

Representations of cross product algebras of Podleś quantum spheres

Konrad Schmüdgen and Elmar Wagner

Fakultät für Mathematik und Informatik

Universität Leipzig, Augustusplatz 10, 04109 Leipzig, Germany

E-mail: Konrad.Schmuedgen@math.uni-leipzig.de / Elmar.Wagner@math.uni-leipzig.de

Abstract

Hilbert space representations of the cross product $*$ -algebras of the Hopf $*$ -algebra $\mathcal{U}_q(\mathfrak{su}_2)$ and its module $*$ -algebras $\mathcal{O}(S^2_{qr})$ of Podleś spheres are investigated and classified by describing the action of generators. The representations are analyzed within two approaches. It is shown that the Hopf $*$ -algebra $\mathcal{O}(\mathrm{SU}_q(2))$ of the quantum group $\mathrm{SU}_q(2)$ decomposes into an orthogonal sum of projective Hopf modules corresponding to irreducible integrable $*$ -representations of the cross product algebras and that each irreducible integrable $*$ -representation appears with multiplicity one. The projections of these projective modules are computed. The decompositions of tensor products of irreducible integrable $*$ -representations with spin l representations of $\mathcal{U}_q(\mathfrak{su}_2)$ are given. The invariant state h on $\mathcal{O}(S^2_{qr})$ is studied in detail. By passing to function algebras over the quantum spheres S^2_{qr} , we give chart descriptions of quantum line bundles and describe the representations from the first approach by means of the second approach.

Keywords: Quantum groups, unbounded representations

Mathematics Subject Classifications (2000): 17B37, 81R50, 46L87

0 Introduction

Podleś quantum spheres [13] (see [7, Section 4.5] for a short treatment) are a one-parameter family S^2_{qr} , $r \in [0, \infty]$, of mutually non-isomorphic quantum homogeneous spaces of the quantum group $\mathrm{SU}_q(2)$, where $0 < q < 1$. Each of

these spaces can be considered as a quantum analogue of the classical 2-sphere. Their coordinate algebras $\mathcal{O}(S_{qr}^2)$ are right coideal $*$ -subalgebras of the coordinate Hopf $*$ -algebra $\mathcal{O}(\mathrm{SU}_q(2))$ of the quantum group $\mathrm{SU}_q(2)$ and hence left module $*$ -algebras of the Hopf $*$ -algebra $\mathcal{U}_q(\mathrm{su}_2)$. Therefore, the left cross product $*$ -algebra $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathrm{su}_2)$ is defined. The subject of this paper are Hilbert space representations of the $*$ -algebra $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathrm{su}_2)$.

There is now an extensive literature on Podleś spheres (see e.g. [1–3, 5, 6, 12]). Let us restate some algebraic results from these papers that are relevant for our investigations. Let $\mathcal{C}$ denote the coalgebra $\mathcal{O}(\mathrm{SU}_q(2))/\mathcal{O}(S_{qr}^2)^+ \mathcal{O}(\mathrm{SU}_q(2))$ with quotient map $\rho : \mathcal{O}(\mathrm{SU}_q(2)) \rightarrow \mathcal{C}$, where $\mathcal{O}(S_{qr}^2)^+ = \{x \in \mathcal{O}(S_{qr}^2); \varepsilon(x) = 0\}$. The algebra $\mathcal{O}(S_{qr}^2)$ is then the algebra of left $\mathcal{C}$ -covariant elements of $\mathcal{O}(\mathrm{SU}_q(2))$. Note that only in the case $r = 0$ the coalgebra $\mathcal{C}$ is a Hopf algebra and the quantum sphere S_{qr}^2 is the quotient by a quantum subgroup. M. Dijkhuizen and T. Koornwinder [3] have found a skew-primitive element $X_r \in \mathcal{U}_q(\mathrm{su}_2)$ such that $\mathcal{O}(S_{qr}^2)$ is the subalgebra of right X_r -invariant elements of $\mathcal{O}(\mathrm{SU}_q(2))$. A major step have been the results of E. F. Müller and H.-J. Schneider [12]. They showed that $\mathcal{O}(\mathrm{SU}_q(2))$ is faithfully flat as a left (and right) $\mathcal{O}(S_{qr}^2)$ -module and that $\mathcal{C}$ is spanned by group-like elements. As a consequence, $\mathcal{C}$ is the direct sum of simple subalgebras $\mathcal{C}_j$. Then

$$M_j = \{x \in \mathcal{O}(\mathrm{SU}_q(2)); \rho(x_{(1)}) \otimes x_{(2)} \in \mathcal{C}_j \otimes \mathcal{O}(\mathrm{SU}_q(2))\}$$

is a finitely generated projective relative $(\mathcal{O}(\mathrm{SU}_q(2)), \mathcal{O}(S_{qr}^2))$ -Hopf module and $\mathcal{O}(\mathrm{SU}_q(2))$ is the direct sum of these Hopf modules M_j [12, p. 186]. In the subgroup case $r = 0$, the corresponding projections and their Chern numbers have been computed in [6] and [5], respectively. A family of group-like elements spanning the coalgebra $\mathcal{C}$ was determined in [2].

In the present paper, we reconsider and extend these algebraic results in the Hilbert space setting. We prove that $\mathcal{O}(\mathrm{SU}_q(2))$ is the orthogonal direct sum of the Hopf modules M_j and we give an explicit description of this decomposition. Here the skew-primitive element X_r plays a crucial role. Moreover, we determine the projections of the projective modules M_j . All this is carried out in Section 3. Each Hopf module M_j corresponds to an irreducible $*$ -representation π_j of the cross product $*$ -algebra $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathrm{su}_2)$ such that the restriction of π_j to $\mathcal{U}_q(\mathrm{su}_2)$ is a direct sum of spin l representations T_l , $l \in \frac{1}{2}\mathbb{N}_0$. Let us call a $*$ -representation of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathrm{su}_2)$ integrable if it has the latter property. In Section 4, we classify integrable $*$ -representations of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathrm{su}_2)$ and prove that each irreducible integrable $*$ -representation of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathrm{su}_2)$ is unitarily equivalent to one of the

representations π_j . In the course of this classification, we describe the structure of these $*$ -representations π_j by explicit formulas for the actions of generators of $\mathcal{O}(S_{qr}^2)$ on an orthonormal basis of weight vectors for the representation T_l of $\mathcal{U}_q(\mathfrak{su}_2)$. In the terminology of our previous paper [15], this is the first approach to representations of the cross product algebra $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$. We also derive a formula for the decomposition of the tensor product representation $\pi_j \otimes T_l$ into a direct sum of representations π_j .

The corresponding second approach [15] is developed in Section 5. Here we begin with a representation of the $*$ -algebra $\mathcal{O}(S_{qr}^2)$ given in a canonical form and we extend it to a representation of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$. The main technical tool for this is to “decouple” the cross relations of the cross product algebra by finding an auxiliary $*$ -subalgebra $\mathcal{Y}_r$ of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ which commutes with the $*$ -algebra $\mathcal{O}(S_{qr}^2)$. Such decouplings have been found and studied in [4].

Section 6 starts by defining algebras of functions which extend the coordinate algebras $\mathcal{O}(S_{qr}^2)$ and by describing invariant functionals on such function algebras. The algebras of functions together with the invariant functionals will be used to give another description of irreducible integrable $*$ -representations. It should be emphasized that though all irreducible integrable representations π_j of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ involve unbounded operators, their restrictions to the $*$ -subalgebra $\mathcal{O}(S_{qr}^2) \otimes \mathcal{Y}_r$ are given by bounded operators only. In Subsection 6.2, we show that the restriction of an irreducible integrable $*$ -representation to $\mathcal{O}(S_{qr}^2) \otimes \mathcal{Y}_r$ decomposes into a direct sum of two representations which can be realized on algebras of functions with support in the positive and the negative spectrum of a certain self-adjoint operator. This self-adjoint operator represents a coordinate function of the quantum sphere and the two algebras of functions can be considered as “charts” of the projective module M_j . The representation of $\mathcal{O}(S_{qr}^2) \otimes \mathcal{Y}_r$ on each chart leads again to an irreducible $*$ -representation of the cross product algebra which is not integrable and can be described by the formulas from the second approach. To round off this circle of investigations, we recover in Section 7 the irreducible integrable representation by taking the direct sum of both charts and passing to another domain.

In Section 1, we briefly mention the correspondence between relative Hopf modules and modules of cross product algebras and we characterize direct sums of Heisenberg representations for cross product algebras $\mathcal{A} \rtimes \mathcal{U}$ of Hopf $*$ -algebras $\mathcal{A}$ of compact quantum groups. Section 2 collects a number of definitions and basic facts on Podleś spheres and on the cross product algebras $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ which are needed in what follows.

All facts and notions on quantum groups used in this paper can be found, for

instance, in [7]. The algebra $\mathcal{U}_q(\mathfrak{sl}_2)$ is due to P. P. Kulish and N. Y. Reshetikhin [8]. The quantum group $\mathrm{SU}_q(2)$ was invented in [19] and [17] and the quantum spheres S_{qr}^2 were discovered in [13].

Let us introduce some notation. Throughout this paper, q is a real number of the open interval $(0, 1)$. We abbreviate

$$\lambda := q - q^{-1}, \quad \lambda_n := (1 - q^{2n})^{1/2}, \quad [n] := (q^n - q^{-n})/(q - q^{-1}),$$

where $n \in \mathbb{N}_0$. For an algebra $\mathcal{X}$, the notation $\mathcal{X}^n$ stands for the direct sum of n copies of $\mathcal{X}$ and $M_n(\mathcal{X})$ denotes the set of $n \times n$ -matrices with entries from $\mathcal{X}$. Let I be an at most countable index set, V a linear space and $\mathcal{D} = \bigoplus_{i \in I} V_i$, where $V_i = V$ for all $i \in I$. We denote by η_i the vector of $\mathcal{D}$ which has the element $\eta \in V$ as its i -th component and zero otherwise. It is understood that $\eta_i = 0$ whenever $i \notin I$. If V is a Hilbert space, then $\bar{\bigoplus}_{i \in I} V_i$ denotes the closed linear span of $\{\eta_i; \eta \in V, i \in I\}$. Given a dense linear subspace $\mathcal{D}$ of a Hilbert space,

$$\mathcal{L}^+(\mathcal{D}) := \{y \in \mathrm{End}(\mathcal{D}); \mathcal{D} \subset \mathcal{D}(y^*), y^*\mathcal{D} \subset \mathcal{D}\}$$

is a unital $*$ -algebra of closeable operators with involution $y \mapsto y^*$ on $\mathcal{D}$. As customary, $\mathcal{D}(x)$ denotes the domain of an operator x . By a $*$ -representation of a unital $*$ -algebra $\mathcal{X}$ on the domain $\mathcal{D}$, we mean a unit preserving $*$ -homomorphism π from $\mathcal{X}$ into $\mathcal{L}^+(\mathcal{D})$ (see e.g. [14]). When no confusion can arise, we omit the letter which denotes the representation and write x instead of $\pi(x)$.

1 Relative Hopf modules and representations of the cross product algebras

Let $\mathcal{U}$ be a Hopf $*$ -algebra and let $\mathcal{X}$ be a left $\mathcal{U}$ -module $*$ -algebra, that is, $\mathcal{X}$ is a unital $*$ -algebra with left $\mathcal{U}$ -action $\triangleright$ satisfying

$$f \triangleright xy = (f_{(1)} \triangleright x)(f_{(2)} \triangleright y), \quad f \triangleright 1 = \varepsilon(f)1, \quad (f \triangleright x)^* = S(f)^* \triangleright x^* \quad (1)$$

for $x, y \in \mathcal{X}$ and $f \in \mathcal{U}$. Here $\Delta(f) = f_{(1)} \otimes f_{(2)}$ is the Sweedler notation for the comultiplication $\Delta(f)$ of $f \in \mathcal{U}$. Then the *left cross product $*$ -algebra* $\mathcal{X} \rtimes \mathcal{U}$ is the $*$ -algebra generated by the two $*$ -subalgebras $\mathcal{X}$ and $\mathcal{U}$ with respect to the cross commutation relations

$$fx = (f_{(1)} \triangleright x)f_{(2)}, \quad x \in \mathcal{X}, \quad f \in \mathcal{U}. \quad (2)$$

Let $\mathcal{A}$ be a Hopf algebra and $\mathcal{X}$ a right $\mathcal{A}$ -coideal subalgebra of $\mathcal{A}$. A relative Hopf module in ${}_{\mathcal{X}}\mathcal{M}^{\mathcal{A}}$ (see e.g. [11]) is a right $\mathcal{A}$ -comodule M which is also a left $\mathcal{X}$ -module such that the right coaction of $\mathcal{A}$ is $\mathcal{X}$ -linear, that is,

$$(xm)_{(1)} \otimes (xm)_{(2)} = x_{(1)}m_{(1)} \otimes x_{(2)}m_{(2)}, \quad x \in \mathcal{X}, \quad m \in M. \quad (3)$$

Here we write simply xm for the left module action of x at m and we use the Sweedler notation for the coaction.

Let $\langle \cdot, \cdot \rangle$ be a dual pairing of Hopf algebras $\mathcal{U}$ and $\mathcal{A}$. Then any right $\mathcal{A}$ -comodule M determines a left $\mathcal{U}$ -module by

$$f \triangleright m := \langle f, m_{(2)} \rangle m_{(1)}, \quad f \in \mathcal{U}, \quad m \in M, \quad (4)$$

and the right $\mathcal{A}$ -comodule algebra $\mathcal{X}$ becomes a left $\mathcal{U}$ -module algebra. Hence the cross product algebra $\mathcal{X} \rtimes \mathcal{U}$ is defined. The cross relations (2) are given by

$$fx = \langle f_{(1)}, x_{(2)} \rangle x_{(1)}f_{(2)}, \quad x \in \mathcal{X}, \quad f \in \mathcal{U}. \quad (5)$$

The following simple well-known lemma is crucial in what follows.

Lemma 1.1 *Let M be a left $\mathcal{X}$ -module and a right $\mathcal{A}$ -comodule.*

- (i) *If $M \in {}_{\mathcal{X}}\mathcal{M}^{\mathcal{A}}$, then M is a left $\mathcal{X} \rtimes \mathcal{U}$ -module.*
- (ii) *Suppose that $\mathcal{U}$ separates the points of $\mathcal{A}$, that is, $\langle f, a \rangle = 0$ for all $f \in \mathcal{U}$ implies $a = 0$. If M is a left $\mathcal{X} \rtimes \mathcal{U}$ -module, then $M \in {}_{\mathcal{X}}\mathcal{M}^{\mathcal{A}}$.*

Proof. As M is a right $\mathcal{A}$ -comodule, it is a left $\mathcal{U}$ -module. From (4), we obtain

$$f \triangleright (xm) = \langle f, (xm)_{(2)} \rangle (xm)_{(1)}, \quad (6)$$

$$\langle f_{(1)}, x_{(2)} \rangle x_{(1)}(f_{(2)} \triangleright m) = \langle f, x_{(2)}m_{(2)} \rangle x_{(1)}m_{(1)}. \quad (7)$$

If $M \in {}_{\mathcal{X}}\mathcal{M}^{\mathcal{A}}$, then (3) holds and hence the right hand sides of (6) and (7) coincide. That is, the cross relations (5) of $\mathcal{X} \rtimes \mathcal{U}$ are satisfied, so we have a well defined left $\mathcal{X} \rtimes \mathcal{U}$ -module. Conversely, if the right hand sides of (6) and (7) are equal, then (3) follows by using the assumption that $\mathcal{U}$ separates $\mathcal{A}$. $\square$

Suppose now that $\langle \cdot, \cdot \rangle$ is a dual pairing of Hopf $*$ -algebras $\mathcal{U}$ and $\mathcal{A}$ and that $\mathcal{X}$ is a $*$ -invariant right $\mathcal{A}$ -coideal. Suppose that there exists a positive linear functional h on $\mathcal{X}$ (i.e. $h(x^*x) \geq 0$ for $x \in \mathcal{X}$) which is $\mathcal{U}$ -invariant (i.e. $h(f \triangleright x) = \varepsilon(f)h(x)$ for $x \in \mathcal{X}$ and $f \in \mathcal{U}$). Then there is a unique $*$ -representation π_h ,

called the *Heisenberg representation* of the $*$ -algebra $\mathcal{A} \rtimes \mathcal{U}$, with domain $\mathcal{D}_h$ such that $\pi_h|_{\mathcal{A}}$ is the GNS-representation of $\mathcal{A}$ with cyclic vector $\varphi_h = \pi_h(1)$, $\mathcal{D}_h = \pi(\mathcal{A})\varphi_h$ and $\pi(f)\varphi_h = \varepsilon(f)\varphi_h$ for $f \in \mathcal{U}$. (Note that we have taken in [15] the closure of π_h as the Heisenberg representation obtained from h .)

Recall that a Hopf $*$ -algebra $\mathcal{A}$ is called a *CQG-algebra* (compact quantum group algebra) if $\mathcal{A}$ is the linear span of matrix elements of finite dimensional unitary corepresentations of $\mathcal{A}$ [3], [7, Subsection 11.3.1]. A CQG-algebra has a unique Haar state h .

The next proposition is a Hilbert space version of the well-known algebraic fact that Hopf modules in ${}_{\mathcal{A}}\mathcal{M}^{\mathcal{A}}$ are trivial.

Proposition 1.2 *Suppose that $\langle \cdot, \cdot \rangle$ is a dual pairing of a Hopf $*$ -algebra $\mathcal{U}$ and a CQG Hopf $*$ -algebra $\mathcal{A}$ such that $\mathcal{U}$ separates the points of $\mathcal{A}$. Let π be a $*$ -representation of the cross product $*$ -algebra $\mathcal{A} \rtimes \mathcal{U}$ on a domain $\mathcal{D}$. Then π is unitarily equivalent to a direct sum of Heisenberg representations π_h of $\mathcal{A} \rtimes \mathcal{U}$ if and only if $\mathcal{D}$ is a right $\mathcal{A}$ -comodule such that $\pi(f)\varphi = f \triangleright \varphi$ for $f \in \mathcal{U}$ and $\varphi \in \mathcal{D}$.*

Proof. To prove the necessity, it suffices to check the above condition for the Heisenberg representation π_h . Because the Haar state of $\mathcal{A}$ is faithful, there is a well defined right coaction ϕ of $\mathcal{A}$ on $\mathcal{D}$ given by $\phi(\varphi) = \pi_h(a_{(1)})\varphi_h \otimes a_{(2)}$ for $\varphi = \pi_h(a)\varphi_h$, $a \in \mathcal{A}$. Let $f \in \mathcal{U}$. Using the cross relation (2) and $\mathcal{U}$ -invariance of the cyclic vector φ_h , we obtain

$$\pi_h(f)\varphi = \pi_h(fa)\varphi_h = \langle f_{(1)}, a_{(2)} \rangle \pi_h(a_{(1)}f_{(2)})\varphi_h = \langle f, a_{(2)} \rangle \pi_h(a_{(1)})\varphi_h = f \triangleright \varphi.$$

Now we prove the sufficiency. Suppose $\mathcal{D}$ is a right $\mathcal{A}$ -comodule such that $\pi(f)\varphi = f \triangleright \varphi \equiv \langle f, \varphi_{(2)} \rangle \varphi_{(1)}$ for $\varphi \in \mathcal{D}$ and $f \in \mathcal{U}$. Let $\mathcal{D}_{\text{inv}}$ be the subspace of vectors $\omega(\varphi) := \pi(S^{-1}(\varphi_{(2)}))\varphi_{(1)}$, $\varphi \in \mathcal{D}$. Let $f \in \mathcal{U}$ and $\varphi \in \mathcal{D}$. Using the fact that π is a representation and relation (2), we compute

$$\begin{aligned} \pi(f)\omega(\varphi) &= \pi(fS^{-1}(\varphi_{(2)}))\varphi_{(1)} = \langle f_{(1)}, S^{-1}(\varphi_{(2)}) \rangle \pi(S^{-1}(\varphi_{(3)}))\pi(f_{(2)})\varphi_{(1)} \\ &= \langle f_{(1)}, S^{-1}(\varphi_{(3)}) \rangle \pi(S^{-1}(\varphi_{(4)}))\langle f_{(2)}, \varphi_{(2)} \rangle \varphi_{(1)} \\ &= \langle f, S^{-1}(\varphi_{(3)})\varphi_{(2)} \rangle \pi(S^{-1}(\varphi_{(4)}))\varphi_{(1)} = \varepsilon(f)\omega(\varphi). \end{aligned}$$

Hence the functional $\langle \pi(\cdot)\omega(\varphi), \omega(\varphi) \rangle$ on $\mathcal{A}$ is $\mathcal{U}$ -invariant. Since $\mathcal{U}$ separates the points of $\mathcal{A}$, it is also $\mathcal{A}$ -invariant and so a multiple of the Haar state h . Thus, we have

$$\langle \pi(x)\omega(\varphi), \omega(\varphi) \rangle = \|\omega(\varphi)\|^2 h(x), \quad x \in \mathcal{A}. \quad (8)$$

Next we show that $\omega(\varphi) \perp \omega(\varphi')$ for $\varphi, \varphi' \in \mathcal{D}$ implies

$$\pi(\mathcal{A} \rtimes \mathcal{U})\omega(\varphi) \perp \pi(\mathcal{A} \rtimes \mathcal{U})\omega(\varphi'). \quad (9)$$

Let $u \in \mathbb{C}$, $|u| = 1$, and let $x \in \mathcal{A}$ be hermitian. Clearly, $\omega(\varphi) + u\omega(\varphi') \in \mathcal{D}_{\text{inv}}$. By (8), we obtain

$$\begin{aligned} (\|\omega(\varphi)\|^2 + \|\omega(\varphi')\|^2)h(x) &= \langle \pi(x)(\omega(\varphi) + u\omega(\varphi')), \omega(\varphi) + u\omega(\varphi') \rangle \\ &= \langle \pi(x)\omega(\varphi), \omega(\varphi) \rangle + \langle \pi(x)\omega(\varphi'), \omega(\varphi') \rangle + 2\text{Re } u \langle \pi(x)\omega(\varphi), \omega(\varphi') \rangle \\ &= (\|\omega(\varphi)\|^2 + \|\omega(\varphi')\|^2)h(x) + 2\text{Re } u \langle \pi(x)\omega(\varphi), \omega(\varphi') \rangle. \end{aligned}$$

Therefore, $\text{Re } u \langle \pi(x)\omega(\varphi), \omega(\varphi') \rangle = 0$ for all $u \in \mathbb{C}$ with $|u| = 1$ which implies $\langle \pi(x)\omega(\varphi), \omega(\varphi') \rangle = 0$. Since $\mathcal{A}$ is spanned by hermitian elements and π is a $*$ -representation, $\pi(\mathcal{A})\omega(\varphi) \perp \pi(\mathcal{A})\omega(\varphi')$. From (2), we obtain $\pi(f)\pi(\mathcal{A})\omega(\psi) \subseteq \pi(\mathcal{A})\omega(\psi)$ for all $\psi \in \mathcal{D}$, so (9) follows.

Using Zorn's lemma, we choose a maximal set $\{\omega(\varphi_i) ; i \in I\}$ of orthonormal vectors of the vector space $\mathcal{D}_{\text{inv}}$. Let $\mathcal{D}_i = \pi(\mathcal{A} \rtimes \mathcal{U})\omega(\varphi_i)$ and let π_i be the restriction of the representation π to $\mathcal{D}_i$. Since $\langle \pi_i(x)\omega(\varphi_i), \omega(\varphi_i) \rangle = h(x)$ for $x \in \mathcal{A}$ by (8), π_i is unitarily equivalent to the Heisenberg representation π of $\mathcal{A} \rtimes \mathcal{U}$. By (9), $\mathcal{D}_i \perp \mathcal{D}_j$ for $i \neq j$. Since $\varphi = \pi(\varphi_{(2)})\omega(\varphi_{(1)})$ for $\varphi \in \mathcal{D}$, the domain $\mathcal{D}$ is the linear span of subspaces $\mathcal{D}_i$, $i \in I$. Putting the preceding together, we have shown that π is unitarily equivalent to a direct sum of Heisenberg representations. $\square$

Let $\mathcal{A}$ be the CQG -algebra $\mathcal{O}(\text{SU}_q(2))$ and let $\mathcal{U} = \mathcal{U}_q(\text{su}_2)$. Then the hypothesis in Proposition 1.2 is satisfied if and only if π is integrable, that is, its restriction to $\mathcal{U}_q(\text{su}_2)$ is a direct sum of spin l representations T_l , $l \in \frac{1}{2}\mathbb{N}_0$. Therefore, by Proposition 1.2, a $*$ -representation π of the $*$ -algebra $\mathcal{O}(\text{SU}_q(2)) \rtimes \mathcal{U}_q(\text{su}_2)$ is a direct sum of Heisenberg representations π_h if and only if π is integrable. This was proved in [15] by another method. A similar result holds for compact forms of standard quantum groups.

2 The cross product algebra $\mathcal{O}(\text{S}_{qr}^2) \rtimes \mathcal{U}_q(\text{su}_2)$

The Hopf $*$ -algebra $\mathcal{U}_q(\text{su}_2)$ is generated by elements E, F, K, K^{-1} with relations

$$KK^{-1} = K^{-1}K = 1, \quad KE = qEK, \quad FK = qKF, \quad EF - FE = \lambda^{-1}(K^2 - K^{-2}), \quad (10)$$

involution $E^* = F, K^* = K$, comultiplication

$$\Delta(E) = E \otimes K + K^{-1} \otimes E, \quad \Delta(F) = F \otimes K + K^{-1} \otimes F, \quad \Delta(K) = K \otimes K,$$

count $\varepsilon(E) = \varepsilon(F) = \varepsilon(1 - K) = 0$ and antipode $S(K) = K^{-1}$, $S(E) = -qE$, $S(F) = -q^{-1}F$. There is a dual pairing $\langle \cdot, \cdot \rangle$ of the Hopf $*$ -algebras $\mathcal{U}_q(\mathfrak{su}_2)$ and $\mathcal{O}(\mathrm{SU}_q(2))$ given on generators by

$$\langle K^{\pm 1}, d \rangle = \langle K^{\mp 1}, a \rangle = q^{\pm 1/2}, \quad \langle E, c \rangle = \langle F, b \rangle = 1$$

and zero otherwise, where a, b, c, d are the usual generators of $\mathcal{O}(\mathrm{SU}_q(2))$ (see e.g. [7, Chapter 4]).

We shall use the definition of the coordinate algebras $\mathcal{O}(S_{qr}^2)$, $r \in [0, \infty]$, of Podleś spheres as given in [13]. For $r \in [0, \infty)$, $\mathcal{O}(S_{qr}^2)$ is the $*$ -algebra with generators $A = A^*, B, B^*$ and defining relations

$$AB = q^{-2}BA, \quad AB^* = q^2B^*A, \quad B^*B = A - A^2 + r, \quad BB^* = q^2A - q^4A^2 + r. \quad (11)$$

For $r = \infty$, the defining relations of $\mathcal{O}(S_{q,\infty}^2)$ are

$$AB = q^{-2}BA, \quad AB^* = q^2B^*A, \quad B^*B = -A^2 + 1, \quad BB^* = -q^4A^2 + 1. \quad (12)$$

Let $r < \infty$. As shown in [13], $\mathcal{O}(S_{qr}^2)$ is a right $\mathcal{O}(\mathrm{SU}_q(2))$ -comodule $*$ -algebra such that

$$x_{-1} := q^{-1}(1 + q^2)^{1/2}B, \quad x_1 := -(1 + q^2)^{1/2}B^*, \quad x_0 := 1 - (1 + q^2)A. \quad (13)$$

transform by the spin 1 matrix corepresentation (t_{ij}^1) of $\mathrm{SU}_q(2)$. Hence $\mathcal{O}(S_{qr}^2)$ is a left $\mathcal{U}_q(\mathfrak{su}_2)$ -module $*$ -algebra with left action given by $f \triangleright x_j = \sum_i x_i \langle f, t_{ij}^1 \rangle$ for $f \in \mathcal{U}_q(\mathfrak{su}_2)$, $j = -1, 0, 1$. Inserting the form of the matrix (t_{ij}^1) (see [13] or [7, Subsection 4.5.1]) and the Hopf algebra pairing $\langle \cdot, \cdot \rangle$ into (5), we derive the following cross relations for the cross product algebra $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$:

$$\begin{aligned} KA &= AK, \quad EA = AE + q^{-1/2}B^*K, \quad FA = AF - q^{-3/2}BK, \\ KB &= q^{-1}BK, \quad EB = qBE - q^{1/2}(1 + q^2)AK + q^{1/2}K, \quad FB = qBF, \\ KB^* &= qB^*K, \quad EB^* = q^{-1}B^*E, \quad FB^* = q^{-1}B^*F + q^{-1/2}(1 + q^2)AK - q^{-1/2}K. \end{aligned}$$

That is, $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ is the $*$ -algebra with generators $A, B, B^*, E, F, K, K^{-1}$, with defining relations (10), (11) and the preceding set of cross relations.

For $r = \infty$, we set

$$x_{-1} := q^{-1}(1 + q^2)^{1/2}B, \quad x_1 := -(1 + q^2)^{1/2}B^*, \quad x_0 := -(1 + q^2)A. \quad (14)$$

Then the cross relations for $\mathcal{O}(S_{q\infty}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ can be written as

$$\begin{aligned} KA &= AK, \quad EA = AE + q^{-1/2}B^*K, \quad FA = AF - q^{-3/2}BK, \\ KB &= q^{-1}BK, \quad EB = qBE - q^{1/2}(1+q^2)AK, \quad FB = qBF, \\ KB^* &= qB^*K, \quad EB^* = q^{-1}B^*E, \quad FB^* = q^{-1}B^*F + q^{-1/2}(1+q^2)AK. \end{aligned}$$

There is an infinitesimal description of Podleś quantum spheres which was discovered in [3]. For $r \in [0, \infty]$, define an element $X_r \in \mathcal{U}_q(\mathfrak{su}_2)$ by

$$\begin{aligned} X_r &= q^{1/2}(q^{-1} - q)^{-1}r^{-1/2}(1 - K^2) + EK + qFK, \quad r \in (0, \infty), \\ X_0 &= 1 - K^2, \quad X_\infty = EK + qFK. \end{aligned} \tag{15}$$

Then $\Delta(X_r) = 1 \otimes X_r + X_r \otimes K^2$. As shown in [3], the right coideal $*$ -subalgebra

$$\mathcal{X} := \{x \in \mathcal{O}(\mathrm{SU}_q(2)) ; \langle X_r, x_{(1)} \rangle x_{(2)} = 0\}$$

of infinitesimal invariants with respect to X_r can be identified with the coordinate $*$ -algebra $\mathcal{O}(S_{qr}^2)$ of Podleś quantum 2-spheres. The generators x_{-1}, x_0, x_1 of $\mathcal{O}(S_{qr}^2)$ are then identified with the elements

$$\begin{aligned} x_{-1} &= (1 + q^{-2})^{1/2}(r^{1/2}a^2 + ac - qr^{1/2}c^2), \\ x_0 &= (1 + q^{-2})r^{1/2}ab + 1 + (q + q^{-1})bc - (1 + q^2)r^{1/2}dc, \\ x_1 &= (1 + q^{-2})^{1/2}(r^{1/2}b^2 + bd - qr^{1/2}d^2) \quad \text{for } r < \infty; \\ x_{-1} &= (1 + q^{-2})^{1/2}(a^2 - qc^2), \\ x_0 &= (1 + q^{-2})(ab - q^2dc), \\ x_1 &= (1 + q^{-2})^{1/2}(b^2 - qd^2) \quad \text{for } r = \infty. \end{aligned}$$

3 Decomposition of $\mathcal{O}(\mathrm{SU}_q(2))$

Throughout this section, we denote by $\mathcal{X}$ the coordinate algebras $\mathcal{O}(S_{qr}^2)$ and by $\mathcal{A}$ and $\mathcal{U}$ the Hopf $*$ -algebras $\mathcal{O}(\mathrm{SU}_q(2))$ and $\mathcal{U}_q(\mathfrak{su}_2)$, respectively.

First we recall a few crucial algebraic results from [12] needed in what follows. Let $\mathcal{C}$ denote the coalgebra $\mathcal{A}/\mathcal{X}^+\mathcal{A}$ with quotient map $\rho : \mathcal{A} \rightarrow \mathcal{C}$, where $\mathcal{X}^+ = \{x \in \mathcal{X} ; \varepsilon(x) = 0\}$. By [12], $\mathcal{C}$ is spanned by group-like elements and, as a consequence, it is the direct sum of simple subcoalgebras $\mathcal{C}_j, j \in \frac{1}{2}\mathbb{Z}$. Set

$$M_j = \{x \in \mathcal{A} ; \rho(x_{(1)}) \otimes x_{(2)} \in \mathcal{C}_j \otimes \mathcal{A}\}.$$

Then M_j is a finitely generated projective Hopf module in ${}_X\mathcal{M}^{\mathcal{A}}$ and $\mathcal{A}$ is the direct sum of all M_j (cf. [12, p. 163]). Since the dual pairing of $\mathcal{U}$ and $\mathcal{A}$ is non-degenerate [7, Section 2.4], it follows from Lemma 1.1 that Hopf modules in ${}_X\mathcal{M}^{\mathcal{A}}$ are in one-to-one correspondence with left $\mathcal{X} \rtimes \mathcal{U}$ -modules.

In this section, we reconsider these algebraic results in the Hilbert space setting. Since the Haar state on $\mathcal{A}$ is faithful, we can equip $\mathcal{A}$ with the inner product

$$\langle a, b \rangle := h(b^*a), \quad a, b \in \mathcal{A}. \quad (16)$$

The Heisenberg representation of $\mathcal{A}$ is just the left $\mathcal{A} \rtimes \mathcal{U}$ -module $\mathcal{A}$ endowed with the inner product (16). Hence each left $\mathcal{X} \rtimes \mathcal{U}$ -module M_j corresponds to a $*$ -representation, denoted by $\hat{\pi}_j$, of the cross product $*$ -algebra $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathrm{su}_2)$. We will return to this representation in Proposition 4.6 below.

In Theorem 3.1, we give a direct proof that $\mathcal{A} = \mathcal{O}(\mathrm{SU}_q(2))$ is the orthogonal direct sum of Hopf modules M_j and describe this decomposition explicitly. Moreover, the projections of the projective $\mathcal{X}$ -modules M_j are computed.

Let us first introduce some more notation. We abbreviate

$$\begin{aligned} \lambda_{\pm} &:= 1/2 \pm (r + 1/4)^{1/2} \text{ for } r < \infty, \quad \lambda_{\pm} := \pm 1 \text{ for } r = \infty, \\ s &:= 0 \text{ for } r = 0, \quad s := -r^{-1/2}\lambda_- \text{ for } r \in (0, \infty), \quad s := 1 \text{ for } r = \infty. \end{aligned}$$

Note that $s = r^{1/2}\lambda_+^{-1}$ when $r \in [0, \infty)$. For $j \in \frac{1}{2}\mathbb{N}$, define

$$u_j = (d + q^{-1}sb)(d + q^{-2}sb) \dots (d + q^{-2j}sb), \quad (17)$$

$$w_j = (a - qsc)(a - q^2sc) \dots (a - q^{2j}sc), \quad (18)$$

$$u_{-j} = E^{2j} \triangleright w_j, \quad (19)$$

and set $u_0 = w_0 = 1$. From [2], we conclude that $\rho_L(u_j), \rho_L(w_j), j \in \frac{1}{2}\mathbb{N}_0$, are group like elements that span the coalgebra $\mathcal{A}/\mathcal{X}^+\mathcal{A}$, where $\rho_L : \mathcal{A} \rightarrow \mathcal{A}/\mathcal{X}^+\mathcal{A}$ is the canonical mapping. (In order to apply the results in [2], one has to interchange the right $\mathcal{A}$ -comodule algebra $\mathcal{X}$ with the left $\mathcal{A}$ -comodule algebra $\theta(\mathcal{X})$ using the $*$ -algebra automorphism and coalgebra anti-homomorphism $\theta : \mathcal{A} \rightarrow \mathcal{A}$ determined by $\theta(a) = a, \theta(d) = d, \theta(b) = -qc, \theta(c) = -q^{-1}b$.) In particular, the simple coalgebras $\mathcal{C}_j$ are given by $\mathcal{C}_j = \mathbb{C}\rho_L(u_j)$ and $\mathcal{C}_{-j} = \mathbb{C}\rho_L(w_j), j \in \frac{1}{2}\mathbb{N}_0$. As a consequence, $u_j \in M_j$ for $j \in \frac{1}{2}\mathbb{Z}$. A crucial role will play the elements

$$v_{kj}^l := N_{kj}^l F^{l-k} \triangleright (x_1^{l-|j|} u_j), \quad l \in \frac{1}{2}\mathbb{N}_0, \quad j, k = -l, -l+1, \dots, l, \quad (20)$$

of $\mathcal{A}$, where $N_{kj}^l = \|F^{l-k} \triangleright (x_1^{l-|j|} u_j)\|^{-1}$. To describe the projection of the projective module M_j , we define a $(2|j|+1) \times (2|j|+1)$ -matrix P_j with entries from $\mathcal{A}$ by

$$P_j = \left(q^{-(n+m)} [2|j|+1]^{-1} v_{nj}^{|j|} v_{mj}^{|j|*} \right)_{n,m=-|j|}^{|j|}.$$

Theorem 3.1 *The decomposition $\mathcal{A} = \bigoplus_{j \in \frac{1}{2}\mathbb{Z}} M_j$ is an orthogonal direct sum with respect to the inner product given by (16). The set $\{v_{kj}^l; l \in \frac{1}{2}\mathbb{N}_0, k, j = -l, -l+1, \dots, l\}$ is an orthonormal basis of $\mathcal{A}$ and*

$$M_j = \text{Lin}\{v_{kj}^{|j|+n}; n \in \mathbb{N}_0, k = -(|j|+n), -(|j|+n)+1, \dots, |j|+n\}.$$

As a left $\mathcal{X}$ -module, M_j is generated by $\{v_{kj}^{|j|}; k = -|j|, -|j|+1, \dots, |j|\}$, the matrices P_j are orthogonal projections in $M_{2|j|+1}(\mathcal{X})$, and M_j is isomorphic to $\mathcal{X}^{2|j|+1} P_j$. Each vector $v_{kj}^{|j|}$ is cyclic for the $$ -representation $\hat{\pi}_j$ of $\mathcal{X} \rtimes \mathcal{U}$ on M_j .*

Proof. Obviously, v_{lj}^l is a highest weight vector of weight l and the linear space $V_j^l := \text{Lin}\{v_{kj}^l; k = -l, -l+1, \dots, l\}$ is an irreducible $\mathcal{U}_q(\mathfrak{su}_2)$ -module of spin l . Hence v_{kj}^l and $v_{k'j'}^{l'}$ are orthogonal whenever $l \neq l'$ or $k \neq k'$ because then they belong to representations of different spin or are vectors of different weights.

It remains to prove that v_{kj}^l and $v_{k'j'}^l$ are orthogonal when $j \neq j'$. The idea of the proof [12] is to show that v_{kj}^l and $v_{k'j'}^l$ are eigenvectors of different eigenvalues of a hermitian operator $\hat{X}_r$ acting on $\mathcal{A}$. Let $\hat{X}_r$ be defined by

$$\hat{X}_r(a) := a \triangleleft X_r = \langle X_r, a_{(1)} \rangle a_{(2)}, \quad a \in \mathcal{A},$$

where $X_r \in \mathcal{U}_q(\mathfrak{su}_2)$ is given by (15). The relation $S(X_r)^* = S(X_r)$ implies that the operator $\hat{X}_r$ is hermitian. Indeed, for $a, b \in \mathcal{A}$, we have

$$\begin{aligned} \langle a, b \triangleleft X_r \rangle &= h((b \triangleleft X_r)^* a) = h((b^* \triangleleft S(X_r)^*) a) = h((b^* \triangleleft S(X_r)) a) \\ &= \varepsilon(X_{r(2)}) h((b^* \triangleleft S(X_{r(1)})) a) = h((b^* \triangleleft S(X_{r(1)}) X_{r(2)})(a \triangleleft X_{r(3)})) \\ &= h(b^*(a \triangleleft X_r)) = \langle a \triangleleft X_r, b \rangle. \end{aligned}$$

For $j \in \frac{1}{2}\mathbb{Z}$, set

$$\begin{aligned} \mu_j &:= 1 - q^{2j} \quad \text{for } r = 0, \\ \mu_j &:= q^{1/2} (q^{-1} - q)^{-1} r^{-1/2} (1 - q^{-2j} \lambda_- - q^{2j} \lambda_+) \quad \text{for } r \in (0, \infty), \\ \mu_j &:= q^{1/2} (q^{-1} - q)^{-1} (q^{-2j} - q^{2j}) \quad \text{for } r = \infty. \end{aligned}$$

We claim that $u_j \triangleleft X_r = \mu_j u_j$, where the vectors u_j , $j \in \frac{1}{2}\mathbb{Z}$, are defined by (17)–(19). Let $r \in (0, \infty)$. For $j = 0$, we have $u_0 \triangleleft X_r = \varepsilon(X_r) = 0$. Assume that the assertion holds for $j \in \frac{1}{2}\mathbb{N}_0$. Using $\Delta(X_r) = 1 \otimes X_r + X_r \otimes K^2$ and $s = -r^{-1/2}\lambda_- = r^{-1/2}\lambda_+^{-1}$, we compute

$$\begin{aligned} u_{j+1/2} \triangleleft X_r &= u_j((d + q^{-(2j+1)}sb) \triangleleft X_r) + u_j \triangleleft X_r((d + q^{-(2j+1)}sb) \triangleleft K^2) \\ &= u_j[(q^{1/2}r^{-1/2}(q^{-1} - q)^{-1}(1 - q) - q^{1/2}r^{-1/2}q^{-2j}\lambda_- + q\mu_j)d \\ &\quad + (q^{1/2}r^{-1/2}(q^{-1} - q)^{-1}(1 - q^{-1}) + q^{1/2}r^{-1/2}q^{2j}\lambda_+ + q^{-1}\mu_j)q^{-(2j+1)}sb] \\ &= \mu_{j+1/2}u_j(d + q^{-(2j+1)}sb) = \mu_{j+1/2}u_{j+1/2}. \end{aligned}$$

By induction, the claim follows for $j \in \frac{1}{2}\mathbb{N}_0$. Similarly, one proves that $w_j \triangleleft X_r = \mu_{-j}w_j$ for $j \in \frac{1}{2}\mathbb{N}$. Since $(E^{2j} \triangleright w_j) \triangleleft X_r = E^{2j} \triangleright (w_j \triangleleft X_r)$, we obtain $u_j \triangleleft X_r = \mu_j u_j$ for all $j \in \frac{1}{2}\mathbb{Z}$. Analogous, but simpler, computations show that the claim also holds for $r = 0$ and $r = \infty$. Using $x_1 \triangleleft X_r = 0$, we obtain

$$\begin{aligned} (N_{kj}^l F^{l-k} \triangleright (x_1^{l-|j|} u_j)) \triangleleft X_r &= N_{kj}^l F^{l-k} \triangleright (x_1^{l-|j|} (u_j \triangleleft X_r) + (x_1^{l-|j|} \triangleleft X_r)(u_j \triangleleft K^2)) \\ &= \mu_j N_{kj}^l F^{l-k} \triangleright (x_1^{l-|j|} u_j), \end{aligned}$$

so that $\hat{X}_r(v_{kj}^l) = \mu_j v_{kj}^l$. Since $\hat{X}_r$ is hermitian and $\mu_j \neq \mu_{j'}$, we conclude that v_{kj}^l and $v_{kj'}^l$ are orthogonal whenever $j \neq j'$.

The decomposition of $\mathcal{A}$ into an orthogonal direct sum is a consequence of above results. Let $t_{kj}^l \in \mathcal{A}$, $l \in \frac{1}{2}\mathbb{N}_0$, $k, j = -l, -l+1, \dots, l$, denote the matrix elements from the Peter-Weyl decomposition of $\mathcal{A}$ [7, Section 4.2]. As v_{kj}^l is a weight vector with weight k of a spin l representation of $\mathcal{U}_q(\mathfrak{su}_2)$, we know that $v_{kj}^l \in \text{Lin}\{t_{ki}^l; i = -l, -l+1, \dots, l\}$. A simple dimension argument shows that $\text{Lin}\{v_{ki}^l; i = -l, -l+1, \dots, l\} = \text{Lin}\{t_{ki}^l; i = -l, -l+1, \dots, l\}$. Since the elements t_{kj}^l span $\mathcal{A}$, we conclude that $\{v_{kj}^l; l \in \frac{1}{2}\mathbb{N}_0, k, j = -l, -l+1, \dots, l\}$ is an orthonormal basis of $\mathcal{A}$. Recall that $v_{|j|,j}^{|j|} = N_{|j|,j}^{|j|} u_j \in M_j$. As $M_j \in \mathcal{X}\mathcal{M}^{\mathcal{A}}$, it follows by the definition of v_{kj}^l that $v_{kj}^l \in M_j$ for all $l = |j|, |j|+1, \dots$ and $k = -l, -l+1, \dots, l$. Since $\mathcal{A}$ is the direct sum of M_j , we conclude that $M_j = \text{Lin}\{v_{k,j}^l; l = |j|, |j|+1, \dots, k = -l, -l+1, \dots, l\}$ and the decomposition $\mathcal{A} = \bigoplus_{j \in \frac{1}{2}\mathbb{Z}} M_j$ is an orthogonal sum.

Writing

$$\begin{aligned} v_{kj}^l &= N_{kj}^l ((F^{l-k})_{(1) \triangleright x_1^{l-|j|}}) (F^{l-k})_{(2) \triangleright u_j} \\ &= N_{kj}^l (N_{|j|,j}^{|j|})^{-1} ((F^{l-k})_{(1) \triangleright x_1^{l-|j|}}) (F^{l-k})_{(2) \triangleright v_{|j|,j}^{|j|}} \end{aligned}$$

and keeping in mind that $(F^{l-k})_{(2) \triangleright v_{|j|,j}^{[j]}} \in \text{Lin}\{v_{kj}^{[j]}; k = -|j|, -|j| + 1, \dots, |j|\}$, it is clear that M_j is generated by $\{v_{kj}^{[j]}; k = -|j|, -|j| + 1, \dots, |j|\}$ as a left $\mathcal{X}$ -module and that $v_{kj}^{[j]}$ is cyclic for the $*$ -representation $\hat{\pi}_j$ of $\mathcal{X} \rtimes \mathcal{U}$ on M_j .

We turn now to the projections of the projective modules M_j . Defining

$$v_j := [2|j| + 1]^{-1/2} (q^{|j|} v_{-|j|,j}^{[j]}, \dots, q^{-|j|} v_{|j|,j}^{[j]})^t, \quad (21)$$

we can write $P_j = v_j v_j^*$. This immediately implies that $P_j^* = P_j$. In order to prove $P_j^2 = P_j$, it is sufficient to show that $v_j^* v_j = 1$. Recall that $K \triangleright v_{kj}^l = q^k v_{kj}^l$ and $F \triangleright v_{kj}^l = [l+k]^{1/2} [l-k+1]^{1/2} v_{k-1,j}^l$ (see also Equation (23) below). By the third equation in (1), $K \triangleright v_{kj}^{l*} = q^{-k} v_{kj}^{l*}$ and $F \triangleright v_{kj}^{l*} = -q^{-1} [l-k]^{1/2} [l+k+1]^{1/2} v_{k+1,j}^{l*}$. From this, we conclude that $v_j^* v_j = [2|j| + 1]^{-1} (q^{2|j|} v_{-|j|,j}^{[j]*} v_{-|j|,j}^{[j]} + \dots + q^{-2|j|} v_{|j|,j}^{[j]*} v_{|j|,j}^{[j]})$ is a linear combination of vectors of weight 0. We show that $v_j^* v_j$ belongs to a spin 0 representation. Since $K \triangleright v_j^* v_j = v_j^* v_j$, this is equivalent to $F \triangleright (v_j^* v_j) = 0$. Inserting the expressions for v_j gives

$$\begin{aligned} F \triangleright (v_j^* v_j) &= \sum_{k=-|j|}^{|j|} q^{-2k} [2|j| + 1]^{-1} \left((F \triangleright v_{kj}^{[j]*}) (K \triangleright v_{kj}^{[j]}) + (K^{-1} \triangleright v_{kj}^{[j]*}) (F \triangleright v_{kj}^{[j]}) \right) \\ &= [2|j| + 1]^{-1} \sum_{k=-|j|}^{|j|} \left(-q^{-k-1} [|j| - k]^{1/2} [|j| + k + 1]^{1/2} v_{k+1,j}^{[j]*} v_{kj}^{[j]} \right. \\ &\quad \left. + q^{-k} [|j| + k]^{1/2} [|j| - k + 1]^{1/2} v_{kj}^{[j]*} v_{k-1,j}^{[j]} \right) \end{aligned}$$

which telescopes to zero. Since $v_j^* v_j$ belongs to a spin 0 representation and $h(v_{kj}^{[j]*} v_{k,j}^{[j]}) = \|v_{k,j}^{[j]}\|^2 = 1$, we have $v_j^* v_j = h(v_j^* v_j) = \sum_{k=-|j|}^{|j|} q^{-2k} [2|j| + 1]^{-1} = 1$ by (21) as desired. Hence P_j is an orthogonal projection.

Next we verify that $v_{nj}^{[j]} v_{mj}^{[j]*}$ belongs to $\mathcal{X}$. In order to do so, we use the fact that $\mathcal{X}$ is the set of elements $x \in \mathcal{A}$ such that $x \triangleleft X_r = 0$. Since $v_{nj}^{[j]} \triangleleft X_r = \hat{X}_r(v_{nj}^{[j]}) = \mu_j v_{nj}^{[j]}$ and $v_{mj}^{[j]*} = N_{mj}^{[j]} S(F^{|j|-m})^* \triangleright u_j^*$, we get

$$\begin{aligned} (v_{nj}^{[j]} v_{mj}^{[j]*}) \triangleleft X_r &= N_{mj}^{[j]} [v_{nj}^{[j]} (S(F^{|j|-m})^* \triangleright u_j^*) \triangleleft X_r + (v_{nj}^{[j]} \triangleleft X_r) (S(F^{|j|-m})^* \triangleright u_j^*) \triangleleft K^2] \\ &= N_{mj}^{[j]} v_{nj}^{[j]} S(F^{|j|-m})^* \triangleright [u_j^* \triangleleft X_r + \mu_j u_j^* \triangleleft K^2]. \end{aligned}$$

Hence it suffices to show that $u_j^* \triangleleft X_r + \mu_j u_j^* \triangleleft K^2 = 0$ for all $j \in \frac{1}{2}\mathbb{Z}$. This can be done by induction. Let $r \in (0, \infty)$. For $j = 0$, the assertion is true since

$u_0^* \triangleleft X_r = 0$ and $\mu_0 = 0$. Assume that $u_j^* \triangleleft X_r + \mu_j u_j^* \triangleleft K^2 = 0$ holds for $j \in \frac{1}{2}\mathbb{N}_0$. Then $u_j^* \triangleleft X_r = -\mu_j(u_j^* \triangleleft K^2)$. As $u_{j+1/2}^* = (a - q^{-2j}sc)u_j^*$, we compute

$$\begin{aligned} u_{j+1/2}^* \triangleleft X_r + \mu_{j+1/2}(u_{j+1/2}^* \triangleleft K^2) &= (a - q^{-2j}sc)(u_j^* \triangleleft X_r) \\ &\quad + (a - q^{-2j}sc) \triangleleft X_r (u_j^* \triangleleft K^2) + \mu_{j+1/2}(a - q^{-2j}sc) \triangleleft K^2 (u_j^* \triangleleft K^2) \\ &= [-\mu_j(a - q^{-2j}sc) + (q^{1/2}r^{-1/2}(q^{-1} - q)^{-1}(1 - q^{-1}) - q^{-1/2}q^{-2j}s)a \\ &\quad - (q^{1/2}r^{-1/2}(q^{-1} - q)^{-1}(1 - q) - q^{2j+3/2}s^{-1})q^{-2j}sc \\ &\quad + \mu_{j+1/2}(q^{-1}a - q^{-2j+1}sc)](u_j^* \triangleleft K^2). \end{aligned}$$

Inserting the expressions for s , μ_j and $\mu_{j+1/2}$, the preceding equation yields zero. This proves that $u_j^* \triangleleft X_r + \mu_j u_j^* \triangleleft K^2 = 0$ and hence $(v_{n_j}^{[j]} v_{m_j}^{[j]*}) \triangleleft X_r = 0$ for $j \in \frac{1}{2}\mathbb{N}_0$. In the same way, one can show that $w_j^* \triangleleft X_r + \mu_{-j} w_j^* \triangleleft K^2 = 0$ for $j \in \frac{1}{2}\mathbb{N}_0$. Since $u_{-j}^* \triangleleft X_r + \mu_{-j} u_{-j}^* \triangleleft K^2 = S(E^{2j})^* \triangleright [w_j^* \triangleleft X_r + \mu_{-j} w_j^* \triangleleft K^2]$, we conclude that $u_j^* \triangleleft X_r + \mu_j u_j^* \triangleleft K^2 = 0$ holds for all $j \in \frac{1}{2}\mathbb{Z}$. For $r = 0$ and $r = \infty$, the proof is similar.

Finally we prove that M_j is isomorphic to $\mathcal{X}^{2|j|+1}P_j$ as a left $\mathcal{X}$ -module. Define a mapping $\Psi_j : \mathcal{X}^{2|j|+1}P_j \rightarrow M_j$ by

$$\Psi_j((y_{-|j|}, \dots, y_{|j|})P_j) := [2|j|+1]^{-1/2} \sum_{k=-|j|}^{|j|} q^{-k} y_k v_{kj}^{[j]}. \quad (22)$$

Recall that $P_j = v_j v_j^*$ and $v_j^* v_j = 1$. Suppose we are given $y_{-|j|}, \dots, y_{|j|} \in \mathcal{X}$ such that $(y_{-|j|}, \dots, y_{|j|})P_j = 0$. Multiplying by v_j from the right yields

$$0 = (y_{-|j|}, \dots, y_{|j|})P_j v_j = (y_{-|j|}, \dots, y_{|j|})v_j(v_j^* v_j) = [2|j|+1]^{-1/2} \sum_{k=-|j|}^{|j|} q^{-k} y_k v_{kj}^{[j]}.$$

Hence Ψ_j is well defined. Furthermore, $[2|j|+1]^{-1/2} \sum_{k=-|j|}^{|j|} q^{-k} y_k v_{kj}^{[j]} = 0$ implies

$$0 = ([2|j|+1]^{-1/2} \sum_{k=-|j|}^{|j|} q^{-k} y_k v_{kj}^{[j]})v_j^* = (y_{-|j|}, \dots, y_{|j|})P_j,$$

so Ψ_j is injective. Since M_j is generated by $\{v_{kj}^{[j]}; k = -|j|, \dots, |j|\}$ as a left $\mathcal{X}$ -module, Ψ_j is also surjective. Whence Ψ_j realizes the desired isomorphism. This completes the proof. $\square$

4 Integrable $*$ -representations of the cross product algebras

4.1 Classification of integrable $*$ -representations of the cross product algebra $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$

For $l \in \frac{1}{2}\mathbb{N}_0$, let T_l denote the type 1 spin l representations of $\mathcal{U}_q(\mathfrak{su}_2)$. Recall that T_l is an irreducible $*$ -representation of the $*$ -algebra $\mathcal{U}_q(\mathfrak{su}_2)$ acting on a $(2l+1)$ -dimensional Hilbert space with orthonormal basis $\{v_j^l; j = -l, -l+1, \dots, l\}$ by the formulas (see, for instance, [7, Subsection 3.2.1])

$$Kv_j^l = q^j v_j^l, \quad Ev_j^l = [l-j]^{1/2}[l+j+1]^{1/2} v_{j+1}^l, \quad Fv_j^l = [l-j+1]^{1/2}[l+j]^{1/2} v_{j-1}^l. \quad (23)$$

As mentioned in the introduction, we call a $*$ -representation of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ *integrable* if it has the following property:

The restriction to $\mathcal{U}_q(\mathfrak{su}_2)$ is the direct sum of representations T_l , $l \in \frac{1}{2}\mathbb{N}_0$.

The aim of this subsection is classify all integrable $*$ -representations of the cross product $*$ -algebra $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$. We shall use the generators x_j defined by (13) (resp. (14)) rather than A, B, B^* .

Suppose we have such a representation acting on the domain $\mathcal{H}$. Then $\mathcal{H}$ can be written as $\mathcal{H} = \bigoplus_{l \in \frac{1}{2}\mathbb{N}_0} V_l$ such that $V_l = \bigoplus_{j=-l}^l V_j^l$, where each V_j^l is the same Hilbert space V_l^l , say, and the generators of $\mathcal{U}_q(\mathfrak{su}_2)$ act on V_l by (23). We claim that there exist operators $\alpha^\pm(l, j) : V_j^l \rightarrow V_{j\pm 1}^l$, $\alpha^0(l, j) : V_j^l \rightarrow V_{j+1}^l$, $\beta^+(l, j) : V_j^l \rightarrow V_{j+1}^{l+1}$, and self-adjoint operators $\beta^0(l, j) : V_j^l \rightarrow V_j^l$ such that

$$x_1 v_j^l = \alpha^+(l, j) v_j^l + \alpha^0(l, j) v_j^l + \alpha^-(l, j) v_j^l, \quad (24)$$

$$x_0 v_j^l = \beta^+(l, j) v_j^l + \beta^0(l, j) v_j^l + \beta^+(l-1, j)^* v_j^l, \quad (25)$$

$$x_{-1} v_j^l = -q^{-1} (\alpha^-(l+1, j-1)^* v_j^l + \alpha^0(l, j-1)^* v_j^l + \alpha^+(l-1, j-1)^* v_j^l) \quad (26)$$

for $v_j^l \in V_j^l$. Indeed, let $v_j^l \in V_j^l$. Since $Kx_1 = qx_1K$, $x_1 v_j^l$ is a weight vector with weight $j+1$, and since $E^{l-j+1} x_1 v_j^l = q^{-l+j-1} x_1 E^{l-j+1} v_j^l = 0$, $x_1 v_j^l$ is in the linear span of vectors $w_{j+1}^r \in V_{j+1}^r$, where $r \leq l+1$. Similarly, replacing x_1 by x_{-1} and E by F , we conclude that $x_{-1} v_j^l$ belongs to the span of vectors $w_{j-1}^r \in V_{j-1}^r$, $r \leq l+1$. Therefore, since $x_{-1} = -q^{-1} x_1^*$, we have $x_{\pm 1} v_j^l \in V_{j\pm 1}^{l-1} \oplus V_{j\pm 1}^l \oplus V_{j\pm 1}^{l+1}$, so $x_{\pm 1} v_j^l$ is of the form (24) (resp. (26)). From the last two relations of (11) (resp. (12)) and from $x_0^* = x_0$, it follows that $x_0 v_j^l$ is of the form (25). Note that all

operators $\alpha^\pm(l, j)$, $\alpha^0(l, j)$, $\beta^+(l, j)$, $\beta^0(l, j)$ are bounded because the operators x_{-1} , x_0 , x_1 are bounded for any $*$ -representation of the $*$ -algebra $\mathcal{O}(S_{qr}^2)$.

Inserting (23) and (24) into the equation $Ex_1v_j^l = q^{-1}x_1Ev_j^l$, we get

$$\begin{aligned} [l-j]^{1/2}[l+j+3]^{1/2}\alpha^+(l, j) &= q^{-1}[l-j]^{1/2}[l+j+1]^{1/2}\alpha^+(l, j+1), \\ [l-j-1]^{1/2}[l+j+2]^{1/2}\alpha^0(l, j) &= q^{-1}[l-j]^{1/2}[l+j+1]^{1/2}\alpha^0(l, j+1), \\ [l-j-2]^{1/2}[l+j+1]^{1/2}\alpha^-(l, j) &= q^{-1}[l-j]^{1/2}[l+j+1]^{1/2}\alpha^-(l, j+1). \end{aligned}$$

The solutions of these recurrence relations are given by

$$\begin{aligned} \alpha^+(l, j) &= q^{-l+j}[l+j+1]^{1/2}[l+j+2]^{1/2}[2l+1]^{-1/2}[2l+2]^{-1/2}\alpha^+(l, l), \\ \alpha^0(l, j) &= q^{-l+j+1}[l-j]^{1/2}[l+j+1]^{1/2}[2l]^{-1/2}\alpha^0(l, l-1), \\ \alpha^-(l, j) &= q^{-l+j+2}[l-j-1]^{1/2}[l-j]^{1/2}[2]^{-1/2}\alpha^-(l, l-2). \end{aligned}$$

Similarly, from the equation $Ex_0v_j^l = (x_0E + [2]^{1/2}x_1K)v_j^l$, we obtain

$$\begin{aligned} [l-j+1]^{1/2}[l+j+2]^{1/2}\beta^+(l, j) &= [l-j]^{1/2}[l+j+1]^{1/2}\beta^+(l, j+1) + [2]^{1/2}q^j\alpha^+(l, j), \\ [l-j]^{1/2}[l+j+1]^{1/2}\beta^0(l, j) &= [l-j]^{1/2}[l+j+1]^{1/2}\beta^0(l, j+1) + [2]^{1/2}q^j\alpha^0(l, j). \end{aligned}$$

The equation $Ex_0v_l^l = (x_0E + [2]^{1/2}x_1K)v_l^l$ yields in addition

$$\beta^+(l, l) = q^l[2]^{1/2}[2l+2]^{-1/2}\alpha^+(l, l). \quad (27)$$

Further, the equation $0 = \langle v_l^l, (x_1F + q[2]^{1/2}x_0K - qFx_1)v_l^l \rangle$ implies that

$$\alpha^0(l, l-1) = -[2]^{1/2}[2l]^{-1/2}q^{l+1}\beta^0(l, l). \quad (28)$$

As a consequence, $\alpha^0(l, l-1)$ is self-adjoint. Using (27) and (28), it follows that the above recurrence relations for $\beta^+(l, j)$ and $\beta^0(l, j)$ have the following solutions:

$$\beta^+(l, j) = q^j[l-j+1]^{1/2}[l+j+1]^{1/2}[2]^{1/2}[2l+1]^{-1/2}[2l+2]^{-1/2}\alpha^+(l, l), \quad (29)$$

$$\beta^0(l, j) = (1 - q^{l+j+1}[l-j][2][2l]^{-1})\beta^0(l, l). \quad (30)$$

From $0 = \langle v_{l-1}^l, (Ex_{-1} - qx_{-1}E - [2]^{1/2}x_0K)v_{l-1}^{l-1} \rangle$, we derive $\beta^+(l-1, l-1) = -q^{-l}[2l-1]^{1/2}\alpha^-(l, l-2)^*$. Combining this equation with (27) gives

$$\alpha^-(l, l-2) = -q^{2l-1}[2]^{1/2}[2l-1]^{-1/2}[2l]^{-1/2}\alpha^+(l-1, l-1)^*. \quad (31)$$

From (27)–(31), it follows that the representation is completely described if the operators $\alpha^+(l, l)$ and $\beta^0(l, l)$ for $l \in \frac{1}{2}\mathbb{N}_0$ are known. Our next aim is to determine these operators. This will be done by induction on l , where consecutive steps increase l by 1.

We begin by analyzing the relations between $\beta^0(l, l)$ and $\alpha^+(l, l)$. Let $r < \infty$ and abbreviate $\rho := 1 + [2]^2 r$. On V_l^l , the two equations $BB^* = r + q^2 A - q^4 A^2$ and $B^*B = r + A - A^2$ lead to the operator relations

$$\begin{aligned} |\alpha^+(l-1, l-1)^*|^2 + \alpha^0(l, l-1)^2 + |\alpha^-(l+1, l-1)^*|^2 \\ = (1 + q^2)^{-1} (q^2 \rho + (1 - q^2) \beta^0(l, l) - (\beta^0(l, l)^2 + |\beta^+(l, l)|^2)), \\ |\alpha^+(l, l)|^2 = (1 + q^2)^{-1} (q^2 \rho - (1 - q^2) q^2 \beta^0(l, l) - q^4 (\beta^0(l, l)^2 + |\beta^+(l, l)|^2)). \end{aligned}$$

Here it is understood that $\alpha^0(0, -1) = 0$ and $\alpha^+(l-1, l-1) = 0$ for $l = 0, 1/2$. Inserting (27), (28) and (31), these two relations give after some calculations

$$\begin{aligned} |\alpha^+(l-1, l-1)^*|^2 + q^{2l+1} [2l+1]^{-1} [2l+2]^{-1} [2l+3] |\alpha^+(l, l)|^2 \\ = [2]^{-1} (q\rho + (q^{-1} - q) \beta^0(l, l) - q^2 [2l]^{-1} ([2l+3] + q^{2l}) \beta^0(l, l)^2), \end{aligned} \quad (32)$$

$$[2l+2]^{-1} [2l+3] |\alpha^+(l, l)|^2 = [2]^{-1} (\rho - (1 - q^2) \beta^0(l, l) - q^2 \beta^0(l, l)^2). \quad (33)$$

Eliminating $\alpha^+(l, l)$ from these equations yields

$$[2][2l+1] |\alpha^+(l-1, l-1)^*|^2 = [2l] \rho + (1 - q^2) [2l+2] \beta^0(l, l) - [2l]^{-1} [2l+2]^2 q^2 \beta^0(l, l)^2. \quad (34)$$

Now we start with the induction procedure. Our first aim is to show that $\beta^0(0, 0) = 0$. A computation similar to the above shows that

$$\begin{aligned} q[2]^{-1} [3] |\alpha^+(0, 0)|^2 &= (1 + q^2)^{-1} (q^2 \rho - (1 - q^2) q^2 \beta^0(0, 0) - q^4 \beta^0(0, 0)^2), \\ q[2]^{-1} [3] |\alpha^+(0, 0)|^2 &= (1 + q^2)^{-1} (q^2 \rho + (1 - q^2) \beta^0(0, 0) - \beta^0(0, 0)^2). \end{aligned}$$

Eliminating $|\alpha^+(0, 0)|^2$ gives $0 = \beta^0(0, 0) - \beta^0(0, 0)^2$. Since $\beta^0(0, 0)$ is self-adjoint, it is an orthogonal projection. Assume to the contrary that $\beta^0(0, 0) \neq 0$. Then there is $v_0^0 \in V_0^0$ such that $\beta^0(0, 0)v_0^0 = v_0^0$ and $\|v_0^0\| = 1$. Note that $h(\cdot) := \langle (\cdot)v_0^0, v_0^0 \rangle$ is the unique $\mathcal{U}_q(\mathfrak{su}_2)$ -invariant state on $\mathcal{O}(S_{qr}^2)$. From this, we conclude $1 = \langle \beta^0(0, 0)v_0^0, v_0^0 \rangle = \langle x_0 v_0^0, v_0^0 \rangle = h(x_0) = 0$, which is a contradiction. Thus $\beta^0(0, 0) = 0$.

For $l = 1/2$, Equation (34) becomes

$$0 = (q[3] \beta^0(1/2, 1/2))^2 - (1 - q^2) [3] \beta^0(1/2, 1/2) - \rho.$$

This operator identity can only be satisfied if the self-adjoint operator $\beta^0(1/2, 1/2)$ has purely discrete spectrum with eigenvalues

$$\beta^0(1/2, 1/2)_{1/2}^\pm := [3]^{-1}(q^{-2}\lambda_\pm - \lambda_\mp).$$

Clearly, $\beta^0(1/2, 1/2)_{1/2}^+ \neq \beta^0(1/2, 1/2)_{1/2}^-$. Denoting the corresponding eigenspaces by $\mathcal{K}_{-1/2}$ and $\mathcal{K}_{1/2}$, we can write $V_{1/2}^{1/2} = \mathcal{K}_{-1/2} \oplus \mathcal{K}_{1/2}$, and $\beta^0(1/2, 1/2)$ acts on $V_{1/2}^{1/2}$ by

$$\begin{aligned}\beta^0(1/2, 1/2)w_{-1/2} &= \beta^0(1/2, 1/2)_{1/2}^- w_{-1/2}, & w_{-1/2} \in \mathcal{K}_{1/2}, \\ \beta^0(1/2, 1/2)w_{1/2} &= \beta^0(1/2, 1/2)_{1/2}^+ w_{1/2}, & w_{1/2} \in \mathcal{K}_{1/2}.\end{aligned}$$

Here we do not exclude the cases $\mathcal{K}_{-1/2} = \{0\}$ and $\mathcal{K}_{1/2} = \{0\}$.

Next let $l \in \frac{1}{2}\mathbb{N}_0$. Assume that there exist (possibly zero) Hilbert spaces $\mathcal{K}_{-l}, \mathcal{K}_{-l+1}, \dots, \mathcal{K}_l$ such that $V_l^l = \mathcal{K}_{-l} \oplus \dots \oplus \mathcal{K}_l$. For $j \in \frac{1}{2}\mathbb{N}_0, j \leq l$, set

$$\beta^0(l, l)_j^\pm = [2l+2]^{-1}([2j](q^{-2}\lambda_\pm - \lambda_\mp) - (1 - q^{-2})[l-j][l+j+1]), \quad (35)$$

$$\begin{aligned}\alpha^+(l, l)_j^\pm &= [2]^{1/2}[2l+3]^{-1/2}[2l+2]^{-1/2}([2l+2]^2(c+1/4) \\ &\quad - \{(q^{-1}-q)[l-j+1][l+j+1]/2 \pm [2j](c+1/4)^{1/2}\}^2)^{1/2}.\end{aligned} \quad (36)$$

where we already inserted (35) into (33). Assume that $\beta^0(l, l)$ acts on V_l^l by

$$\beta^0(l, l)w_{-j} = \beta^0(l, l)_j^- w_{-j}, \quad w_{-j} \in \mathcal{K}_{-j}, \quad \beta^0(l, l)w_j = \beta^0(l, l)_j^+ w_j, \quad w_j \in \mathcal{K}_j. \quad (37)$$

We show that there exist Hilbert spaces $\mathcal{K}_{-(l+1)}$ and $\mathcal{K}_{l+1}$ such that, up to unitary equivalence, $V_{l+1}^{l+1} = \mathcal{K}_{-(l+1)} \oplus \mathcal{K}_{-l} \oplus \dots \oplus \mathcal{K}_l \oplus \mathcal{K}_{l+1}$, the operator $\beta^0(l+1, l+1)$ acts on V_{l+1}^{l+1} by the formulas (37), and $\alpha^+(l, l) : V_l^l \rightarrow V_{l+1}^{l+1}$ is given by

$$\alpha^+(l, l)w_{-j} = \alpha^+(l, l)_j^- w_{-j}, \quad w_{-j} \in \mathcal{K}_{-j}, \quad \alpha^+(l, l)w_j = \alpha^+(l, l)_j^+ w_j, \quad w_j \in \mathcal{K}_j, \quad (38)$$

where $j = -l, -l+1, \dots, l$.

Observe that $\ker B = \{0\}$ (cf. Subsection 5.3 below). Hence $\ker \alpha^+(l, l) = \{0\}$. Let $\alpha^+(l, l) = U|\alpha^+(l, l)|$ denote the polar decomposition of $\alpha^+(l, l)$. Then U is an isometry from V_l^l onto $\alpha^+(l, l)V_l^l$. On the other hand, we have the decomposition $V_{l+1}^{l+1} = \alpha^+(l, l)V_l^l \oplus \ker \alpha^+(l, l)^*$. After applying a unitary transformation, we can assume that $V_{l+1}^{l+1} = V_l^l \oplus \ker \alpha^+(l, l)^* = \mathcal{K}_{-l} \oplus \dots \oplus \mathcal{K}_l \oplus \ker \alpha^+(l, l)^*$ and $\alpha^+(l, l)v_l^l = |\alpha^+(l, l)|v_l^l$ for all $v_l^l \in V_l^l$. From (33) and (37), it follows that the

action of $\alpha^+(l, l)$ on V_l^l is determined by Equation (38) as asserted. We proceed by describing the action of $\beta^0(l+1, l+1)$. By (34), $|\alpha^+(l, l)^*|^2$ commutes with $\beta^0(l+1, l+1)$. Hence $\beta^0(l+1, l+1)$ leaves the subspaces V_l^l and $\ker \alpha^+(l, l)^*$ of V_{l+1}^{l+1} invariant. Let $\tilde{\beta}^0(l+1, l+1)$ and $\hat{\beta}^0(l+1, l+1)$ denote the restrictions of $\beta^0(l+1, l+1)$ to V_l^l and $\ker \alpha^+(l, l)^*$, respectively. Evaluating the relation $0 = \langle (x_0 x_1 - q^2 x_1 x_0 - (1 - q^2) x_1) v_l^l, v_{l+1}^{l+1} \rangle$ for $v_l^l \in V_l^l$ and $v_{l+1}^{l+1} \in V_{l+1}^{l+1}$ yields

$$\begin{aligned} \beta^0(l+1, l+1) \alpha^+(l, l) - q^2 \alpha^0(l+1, l+1) \beta^+(l, l) \\ - q^2 \alpha^+(l, l) \beta^0(l, l) - (1 - q^2) \alpha^+(l, l) = 0. \end{aligned}$$

Inserting (27) and observing that $\alpha^+(l, l) \beta^0(l, l) = \beta^0(l, l) \alpha^+(l, l)$ by (37) and (38), we deduce the following operator equation on V_l^l

$$\{(1 + q^{2l+4}[2][2l+2]^{-1}) \tilde{\beta}^0(l+1, l+1) - q^2 \beta^0(l, l) - (1 - q^2)\} \alpha^+(l, l) = 0.$$

Since $V_l^l = \overline{\alpha^+(l, l) V_l^l}$, the operator in braces must be zero. This implies that

$$\tilde{\beta}^0(l+1, l+1) = [2l+4]^{-1}[2l+2](\beta^0(l, l) + q^{-2} - 1)$$

Hence the operator $\beta^0(l+1, l+1)$ acts on $V_l^l = \mathcal{K}_{-l} \oplus \dots \oplus \mathcal{K}_l$ by

$$\begin{aligned} \beta^0(l+1, l+1) w_j &= [2l+4]^{-1}[2l+2](\beta^0(l, l)_j^\epsilon + q^{-2} - 1) w_j \\ &= [2l+4]^{-1} ([2j](q^{-2} \lambda_\epsilon - \lambda_{-\epsilon} + (q^{-2} - 1)[l-j+1][l+j+2]) w_j, \end{aligned}$$

where $\epsilon = \text{sign}(j)$. The last equation is obtained by straightforward computations. On $\ker \alpha^+(l, l)^*$, Equation (34) reads

$$0 = \rho + (1 - q^2)[2l+2]^{-1}[2l+4] \hat{\beta}^0(l+1, l+1) - ([2l+2]^{-1}[2l+4] q \hat{\beta}^0(l+1, l+1))^2.$$

Since the solution of the quadratic equation $0 = -\rho - (q^{-1} - q)t + t^2$ is given by $t^\pm = q^{-1} \lambda_\pm - q \lambda_\mp$, we conclude that $\hat{\beta}^0(l+1, l+1)$ has purely discrete spectrum consisting of the two distinct eigenvalues

$$\beta^0(l+1, l+1)_{l+1}^\pm = [2l+4]^{-1}[2l+2](q^{-2} \lambda_\pm - \lambda_\mp),$$

and $\ker \alpha^+(l, l)^*$ splits into the direct sum $\ker \alpha^+(l, l)^* = \mathcal{K}_{-(l+1)} \oplus \mathcal{K}_{l+1}$, where $\mathcal{K}_{-(l+1)}$ and $\mathcal{K}_{l+1}$ denote the eigenspaces of $\hat{\beta}^0(l+1, l+1)$ corresponding to the eigenvalues $\beta^0(l+1, l+1)_{l+1}^-$ and $\beta^0(l+1, l+1)_{l+1}^+$, respectively. Accordingly, the operator $\beta^0(l+1, l+1)$ acts on $\ker \alpha^+(l, l)^*$ by

$$\begin{aligned} \beta^0(l+1, l+1) w_{-(l+1)} &= \beta^0(l+1, l+1)_{l+1}^- w_{-(l+1)}, \quad w_{-(l+1)} \in \mathcal{K}_{-(l+1)}, \\ \beta^0(l+1, l+1) w_{l+1} &= \beta^0(l+1, l+1)_{l+1}^+ w_{l+1}, \quad w_{l+1} \in \mathcal{K}_{l+1}. \end{aligned}$$

This shows that $\beta^0(l+1, l+1)$ is of the same form as $\beta^0(l, l)$. By induction, the operators $\beta^0(l, l)$ and $\alpha^+(l, l)$ are now completely determined.

In the case $r = \infty$, there are only minor changes in the preceding reasoning. Set $\rho = (q + q^{-1})^2$. Then, Equations (33)–(34) remain valid if one omits the first order term of $\beta^0(l, l)$. Equations (35) and (36) become

$$\beta^0(l, l)_j^\pm = \pm q^{-1} [2][2j][2l+2]^{-1}, \quad (39)$$

$$\alpha^+(l, l)_j^\pm = [2]^{1/2} [2(l+j+1)]^{1/2} [2(l-j+1)]^{1/2} [2l+2]^{-1/2} [2l+3]^{-1/2}, \quad (40)$$

and $\alpha^+(l, l)$ is again given by (38).

Note that whenever a Hilbert space $\mathcal{K}_j$ is non-zero, it appears as a direct summand in each V_k^l , where $l = |j|, |j| + 1, \dots$ and $k = -l, -l + 1, \dots, l$. Moreover, the generators of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ leave this decomposition invariant. Hence the representation of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ on $\bigoplus_{l \in \frac{1}{2}\mathbb{N}_0} \bigoplus_{k=-l}^l V_k^l$ splits into a direct sum of representations on $\bigoplus_{n \in \mathbb{N}_0} \bigoplus_{k=-(|j|+n)}^{|j|+n} \mathcal{K}_k^{|j|+n}$, where each $\mathcal{K}_k^{|j|+n}$ is the same Hilbert space $\mathcal{K}_j$ and is considered as a direct summand of $V_k^{|j|+n}$.

It still remains to prove that Equations (23)–(26) define a representation of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ when we insert the expressions for the operators obtained in the previous discussion. This can be done by showing that the defining relations of $\mathcal{O}(S_{qr}^2)$ and the cross relations of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ are satisfied. We have checked this; the details of these lengthy and tedious computations are omitted.

We summarize the outcome of above considerations in the next theorem.

Theorem 4.1 *Each integrable $*$ -representation of the cross product $*$ -algebra $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ is, up to unitary equivalence, a direct sum of representations $\pi_j^\pm$, $j \in \frac{1}{2}\mathbb{N}_0$, of the following form:*

The domain is the direct sum $\bigoplus_{l \in \mathbb{N}_0} \bigoplus_{k=-(j+l)}^{j+l} \mathcal{K}_k^{j+l}$, where each $\mathcal{K}_k^{j+l}$ is the same Hilbert space $\mathcal{K}$. The generators E, F, K of $\mathcal{U}_q(\mathfrak{su}_2)$ act on $\bigoplus_{k=-(j+l)}^{j+l} \mathcal{K}_k^{j+l}$ by (23). The actions of the generators x_1, x_0, x_{-1} of $\mathcal{O}(S_{qr}^2)$ are determined by

$$\begin{aligned} x_1 v_k^l &= q^{-l+k} [l+k+1]^{1/2} [l+k+2]^{1/2} [2l+1]^{-1/2} [2l+2]^{-1/2} \alpha^+(l, l)_j^\pm v_{k+1}^{l+1} \\ &\quad - q^{k+2} [l-k]^{1/2} [l+k+1]^{1/2} [2]^{1/2} [2l]^{-1} \beta^0(l, l)_j^\pm v_{k+1}^l \\ &\quad - q^{l+k+1} [l-k-1]^{1/2} [l-k]^{1/2} [2l-1]^{-1/2} [2l]^{-1/2} \alpha^+(l-1, l-1)_j^\pm v_{k+1}^{l-1}, \\ x_0 v_k^l &= q^k [l-k+1]^{1/2} [l+k+1]^{1/2} [2]^{1/2} [2l+1]^{-1/2} [2l+2]^{-1/2} \alpha^+(l, l)_j^\pm v_k^{l+1} \\ &\quad + (1 - q^{l+k+1} [l-k] [2] [2l]^{-1}) \beta^0(l, l)_j^\pm v_k^l \\ &\quad + q^k [l-k]^{1/2} [l+k]^{1/2} [2]^{1/2} [2l-1]^{-1/2} [2l]^{-1/2} \alpha^+(l-1, l-1)_j^\pm v_k^{l-1}, \end{aligned}$$

$$\begin{aligned}
x_{-1}v_k^l &= q^{l+k}[l-k+1]^{1/2}[l-k+2]^{1/2}[2l+1]^{-1/2}[2l+2]^{-1/2}\alpha^+(l, l)_j^\pm v_{k-1}^{l+1} \\
&\quad + q^k[l-k+1]^{1/2}[l+k]^{1/2}[2]^{1/2}[2l]^{-1}\beta^0(l, l)_j^\pm v_{k-1}^l \\
&\quad - q^{-l+k-1}[l+k-1]^{1/2}[l+k]^{1/2}[2l-1]^{-1/2}[2l]^{-1/2}\alpha^+(l-1, l-1)_j^\pm v_{k-1}^{l-1},
\end{aligned}$$

where, for $r < \infty$, the real numbers $\beta^0(l, l)_j^\pm$ and $\alpha^+(l, l)_j^\pm$ are given by (35) and (36), respectively, and, for $r = \infty$, by (39) and (40), respectively.

Representations corresponding to different pairs of labels $(j, \pm)$ are not unitarily equivalent (with only one obvious exception: $\pi_0^+ = \pi_0^-$). A representation of this list is irreducible if and only if $\mathcal{K} = \mathbb{C}$.

An immediate consequence of Theorem 4.1 is the following

Corollary 4.2 (i) Each integrable $*$ -representation of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ is a direct sum of integrable irreducible $*$ -representations.

(ii) Each integrable irreducible $*$ -representation of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ is unitarily equivalent to a $*$ -representations $\pi_j^\pm$, $j \in \frac{1}{2}\mathbb{N}_0$, with $\mathcal{K} = \mathbb{C}$.

It will be convenient to introduce the following notation. If $\pi_j^\pm$, $j \in \frac{1}{2}\mathbb{N}_0$, denotes an irreducible integrable $*$ -representation of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ (that is, $\pi_j^\pm$ is the representation from Theorem 4.1 with $\mathcal{K} = \mathbb{C}$), then we set

$$\pi_j := \pi_{-j}^- \text{ for } j < 0, \quad \pi_j := \pi_j^+ \text{ for } j \geq 0, \quad j \in \frac{1}{2}\mathbb{Z}.$$

4.2 Decomposition of tensor products of irreducible integrable representations with spin l representations

Lemma 4.3 Suppose that $\mathcal{X}$ is a left module $*$ -algebra of a Hopf $*$ -algebra $\mathcal{U}$. Let π and T be $*$ -representations of the $*$ -algebras $\mathcal{X} \rtimes \mathcal{U}$ and $\mathcal{U}$ on domains $\mathcal{D}$ and V , respectively. Then there is a $*$ -representation, denoted by $\pi \otimes T$, of the $*$ -algebra $\mathcal{X} \rtimes \mathcal{U}$ on the domain $\mathcal{D} \otimes V$ such that $(\pi \otimes T)(x) = \pi(x) \otimes T(1)$ for $x \in \mathcal{X}$ and $(\pi \otimes T)(f) = \pi(f_{(1)}) \otimes T(f_{(2)})$ for $f \in \mathcal{U}$.

Proof. It suffices to check that $\pi \otimes T$ respects the cross relation (2), that is, $(\pi \otimes T)(f)(\pi \otimes T)(x) = (\pi \otimes T)(f_{(1)} \triangleright x)(\pi \otimes T)(f_{(2)})$ for $x \in \mathcal{X}$ and $f \in \mathcal{U}$. The details of this easy verification are left to the reader. $\square$

Clearly, the tensor product $\pi \otimes T_l$ of an integrable $*$ -representation π of the cross product algebra $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ and a spin l representation T_l of $\mathcal{U}_q(\mathfrak{su}_2)$ is

again integrable, so Corollary 4.2 applies. The decomposition of $\pi_j \otimes T_l$ into a direct sum of irreducible integrable representations of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ is described in the next proposition.

Proposition 4.4 *For $j \in \frac{1}{2}\mathbb{Z}$ and $l \in \frac{1}{2}\mathbb{N}_0$, let π_j be an irreducible integrable $*$ -representation of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ and T_l a spin l representation of $\mathcal{U}_q(\mathfrak{su}_2)$. Then, up to unitary equivalence, $\pi_j \otimes T_l = \pi_{j-l} \oplus \pi_{j-l+1} \oplus \dots \oplus \pi_{j+l}$.*

Proof. For $l = 0$, there is nothing to prove. Let $l = 1/2$. From Theorem 4.1, it follows that the restriction of π_j to $\mathcal{U}_q(\mathfrak{su}_2)$ is of the form $\oplus_{n \in \mathbb{N}_0} T_{|j|+n}$. Assume first that $j \neq 0$. By the Clebsch–Gordon decomposition, we have

$$\pi_j \otimes T_{1/2} = T_{|j|-1/2} \oplus (\oplus_{n \in \mathbb{N}_0} 2T_{|j|+n+1/2}) \quad (41)$$

as representations of $\mathcal{U}_q(\mathfrak{su}_2)$. By Corollary 4.2, $\pi_j \otimes T_{1/2}$ decomposes into a direct sum of irreducible integrable $*$ -representations described in Theorem 4.1. In (41), there occur no spin k representations for $k < |j| - 1/2$, exactly one spin $|j| - 1/2$ representation, and two spin k representations for $k = |j| + 1/2, |j| + 3/2, \dots$. Thus we must have $\pi_j \otimes T_{1/2} = \pi_{|j|-1/2}^{\epsilon_1} \oplus \pi_{|j|+1/2}^{\epsilon_2}$, where $\epsilon_1, \epsilon_2 \in \{+, -\}$ are to be determined. Moreover, $\pi_{|j|-1/2}^{\epsilon_1}$ and $\pi_{|j|+1/2}^{\epsilon_2}$ are irreducible. Again by the Clebsch–Gordon decomposition,

$$v_{|j|-1/2}^{|j|-1/2} = [2|j|+1]^{-1/2} (q^{1/2}[2|j|]^{1/2} v_{|j|}^{|j|} \otimes v_{-1/2}^{1/2} - q^{-|j|} v_{|j|-1}^{|j|} \otimes v_{1/2}^{1/2}) \quad (42)$$

is the (unique) highest weight vector of the spin $|j| - 1/2$ representation in (41). If $|j| = 1/2$, then v_0^0 belongs to a spin 0 representation and $\pi_{|j|-1/2}^{\epsilon_1} = \pi_0$ is the Heisenberg representation. Let $|j| > 1/2$ and $\epsilon = \text{sign}(j)$. Then a straightforward computation gives

$$\langle v_{|j|-1/2}^{|j|-1/2}, \pi_j \otimes T_{1/2}(x_0) v_{|j|-1/2}^{|j|-1/2} \rangle = [2|j|+2][2|j|-1][2|j|+1]^{-1}[2|j|]^{-1} \beta^0(|j|, |j|)_{|j|}^{\epsilon}.$$

Hence $\epsilon_1 = \text{sign}(\langle v_{|j|-1/2}^{|j|-1/2}, \pi_j \otimes T_{1/2}(x_0) v_{|j|-1/2}^{|j|-1/2} \rangle) = \text{sign}(\beta^0(|j|, |j|)_{|j|}^{\epsilon}) = \text{sign}(j)$. The linear space of highest weight vectors belonging to spin $|j| + 1/2$ representations in (41) is 2-dimensional and spanned by the orthonormal vectors

$$\begin{aligned} u_{|j|+1/2}^{|j|+1/2} &= [2|j|+3]^{-1/2} (q^{1/2}[2|j|+2]^{1/2} v_{|j|+1}^{|j|+1} \otimes v_{-1/2}^{1/2} - q^{-|j|-1} v_{|j|}^{|j|+1} \otimes v_{1/2}^{1/2}), \\ w_{|j|+1/2}^{|j|+1/2} &= v_{|j|}^{|j|} \otimes v_{1/2}^{1/2}. \end{aligned}$$

The vector

$$v := [2|j|+3]^{1/2}[2|j|+2]^{-1/2}\alpha^+ (|j|, |j|)_{|j|}^\epsilon w_{|j|+1/2}^{|j|+1/2} \\ - q[2]^{1/2}[2|j|]^{-1/2}\beta^0 (|j|, |j|)_{|j|}^\epsilon u_{|j|+1/2}^{|j|+1/2}$$

is orthogonal to $x_1 v_{|j|-1/2}^{|j|-1/2}$. Hence $\|v\|^{-1}v$ is the (unique) highest weight vector of the spin $|j|+1/2$ representation belonging to the decomposition of $\pi_{|j|+1/2}^{\epsilon_2}$. Our goal is to determine $\epsilon_2 = \text{sign}(\langle \|v\|^{-1}v, x_0 \|v\|^{-1}v \rangle) = \text{sign}(\langle v, x_0 v \rangle)$. Using (35), (36) and the formulas in Theorem 4.1, we obtain

$$\langle v, x_0 v \rangle = \{ [2|j|+3][2|j|-1][2|j|]^{-1}[2|j|+1]^{-1}(\alpha^+ (|j|, |j|)_{|j|}^\epsilon)^2 + \\ [2][2|j|]^{-1}[2|j|+2]^{-1} \} \beta^0 (|j|, |j|)_{|j|}^\epsilon.$$

As the expression in braces is positive, we deduce $\epsilon_2 = \text{sign}(\beta^0 (|j|, |j|)_{|j|}^\epsilon) = \text{sign}(j)$.

Next, assume that $j = 0$. By similar arguments as above, we conclude that $\pi_0 \otimes T_{1/2} = \pi_{1/2}^{\epsilon_1} \oplus \pi_{1/2}^{\epsilon_2}$. Let u_1 and u_2 denote the highest weight vectors of the spin $1/2$ representations belonging to $\pi_{1/2}^{\epsilon_1}$ and $\pi_{1/2}^{\epsilon_2}$, respectively. Then $\langle u_1, x_0 u_1 \rangle = \beta^0(1/2, 1/2)_{1/2}^{\epsilon_1}$ and $\langle u_2, x_0 u_2 \rangle = \beta^0(1/2, 1/2)_{1/2}^{\epsilon_2}$. The vector $w_{1/2}^{1/2} = v_0^0 \otimes v_{1/2}^{1/2}$ belongs to the span of the orthonormal vectors u_1 and u_2 . Since $\langle w_{1/2}^{1/2}, x_0 w_{1/2}^{1/2} \rangle = 0$, we get $\text{sign}(\beta^0(1/2, 1/2)_{1/2}^{\epsilon_1}) \neq \text{sign}(\beta^0(1/2, 1/2)_{1/2}^{\epsilon_2})$, whence $\epsilon_1 \neq \epsilon_2$. Summarizing the preceding results, we conclude that, for all $j \in \frac{1}{2}\mathbb{Z}$, the representation $\pi_j \otimes T_{1/2}$ decomposes into $\pi_j \otimes T_{1/2} = \pi_{j-1/2} \oplus \pi_{j+1/2}$.

The proposition can now be proved by induction. Let $k \in \frac{1}{2}\mathbb{N}$. Assume that Proposition 4.4 holds for all $l = 0, 1/2, \dots, k$. By Corollary 4.2, $\pi_j \otimes T_k \otimes T_{1/2} = (\pi_j \otimes T_k) \otimes T_{1/2} = \pi_j \otimes (T_k \otimes T_{1/2})$ decomposes into irreducible integrable $*$ -representations. By our induction hypothesis and (41), we have

$$(\pi_j \otimes T_k) \otimes T_{1/2} = \pi_{j-k-1/2} \oplus (\oplus_{n=0}^{2k-1} 2\pi_{j-k+n+1/2}) \oplus \pi_{j+k+1/2}.$$

On the other hand,

$$\pi_j \otimes (T_k \otimes T_{1/2}) = (\oplus_{n=0}^{2k-1} \pi_{j-k+n+1/2}) \oplus (\pi_j \otimes T_{k+1/2}).$$

Comparing both results shows that

$$\pi_j \otimes T_{k+1/2} = \pi_{j-k-1/2} \oplus \pi_{j-k+1/2} \oplus \dots \oplus \pi_{j+k+1/2}. \quad \square$$

4.3 Realization of irreducible integrable $*$ -representations of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ on $\mathcal{O}(\mathrm{SU}_q(2))$

In this subsection, we relate the irreducible representation π_j from Theorem 4.1 to the representation $\hat{\pi}_j$ corresponding to the Hopf module M_j from Section 3.

Lemma 4.5 *Let $l \in \frac{1}{2}\mathbb{N}$. Define $e_{1/2}^+ = d + q^{-2l-1}sb$ and $e_{-1/2}^- = c - q^{2l+1}sa$. Set $e_{-1/2}^+ = F \triangleright e_{1/2}^+$ and $e_{1/2}^- = E \triangleright e_{-1/2}^-$. Then*

$$q^{1/2}[2l]^{1/2}v_{ll}^l e_{-1/2}^+ - q^{-l}v_{l-1,l}^l e_{1/2}^+ = q^{-1/2}[2l]^{1/2}v_{-l,-l}^l e_{1/2}^- - q^l v_{-l+1,-l}^l e_{-1/2}^- = 0. \quad (43)$$

Proof. Recall that $v_{ll}^l = N_{ll}^l u_l = N_{ll}^l (d + q^{-1}sb) \cdot \dots \cdot (d + q^{-2l}sb)$ and that $v_{l-1,l}^l = [2l]^{-1/2} F \triangleright v_{ll}^l = [2l]^{-1/2} N_{ll}^l (F \triangleright u_l)$. A straightforward induction argument shows that $F \triangleright u_l = q^{l-1/2} [2l] u_{l-1/2} (c + q^{-2l}sa)$. Now a direct calculation gives

$$\begin{aligned} u_l e_{-1/2}^+ - q^{-l-1/2} [2l]^{-1} (F \triangleright u_l) e_{1/2}^+ \\ = u_{l-1/2} (d + q^{-2l}sb) (c + q^{-2l-1}sa) - q^{-1} u_{l-1/2} (c + q^{-2l}sa) (d + q^{-2l-1}sb) = 0 \end{aligned}$$

which implies the first equality in (43). The second equality is proved in the same way by using $v_{-l,-l}^l = \|w_l\|^{-1} w_l$. $\square$

Proposition 4.6 *Let $j \in \frac{1}{2}\mathbb{Z}$. The irreducible integrable $*$ -representation π_j of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ from Theorem 4.1 is unitarily equivalent to the $*$ -representation $\hat{\pi}_j$ on the left $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ -module M_j from Theorem 3.1.*

Proof. By Theorem 3.1, the restriction of the representation $\hat{\pi}_j$ to $\mathcal{U}_q(\mathfrak{su}_2)$ is of the form $\oplus_{n \in \mathbb{N}_0} T_{|j|+n}$. Likewise, by Theorem 4.1, the restriction of the representation π_j to $\mathcal{U}_q(\mathfrak{su}_2)$ is $\oplus_{n \in \mathbb{N}_0} T_{|j|+n}$. Therefore, by Corollary 4.2(i), $\hat{\pi}_j$ is unitarily equivalent either to $\pi_{|j|}^-$ or to $\pi_{|j|}^+$. That is, it only remains to specify the label $+$ or $-$. Recall from Theorem 3.1 that M_j is the linear span of vectors $v_{kj}^{|j|+n}$, where $n \in \mathbb{N}_0$ and $k = -(|j| + n), \dots, |j| + n$.

For $j = 0$, there is only the unique Heisenberg representation. A direct calculation yields $\langle v_{1/2,-1/2}^{1/2}, x_0 v_{1/2,-1/2}^{1/2} \rangle = \beta^0(1/2, 1/2)_{1/2}^-$ and $\langle v_{1/2,1/2}^{1/2}, x_0 v_{1/2,1/2}^{1/2} \rangle = \beta^0(1/2, 1/2)_{1/2}^+$ so that $\hat{\pi}_{-1/2} = \pi_{-1/2}$ and $\hat{\pi}_{1/2} = \pi_{1/2}$.

We proceed by induction. Let $l \in \frac{1}{2}\mathbb{N}$. Assume that π_l is unitarily equivalent to $\hat{\pi}_l$. Recall that $T_{1/2}$ is the spin $1/2$ representation on $V_{1/2} = \mathrm{Lin}\{v_{-1/2}^{1/2}, v_{1/2}^{1/2}\}$,

where $v_{\pm 1/2}^{1/2}$ are the weight vectors. By Proposition 4.4, the tensor product representation $\hat{\pi}_l \otimes T_{1/2} \cong \pi_l \otimes T_{1/2}$ decomposes into the direct sum $\pi_{l-1/2} \oplus \pi_{l+1/2}$ on $M_l \otimes V_{1/2} = W_{l-1/2} \oplus W_{l+1/2}$. Set $e_{1/2} := \|d + q^{-2l-1}sb\|^{-1}(d + q^{-2l-1}sb)$ and $e_{-1/2} := F \triangleright e_{1/2}$. Consider the linear mapping $\Phi : M_l \otimes V_{1/2} \rightarrow \mathcal{O}(\mathrm{SU}_q(2))$ defined by $\Phi(v \otimes v_{\pm 1/2}^{1/2}) = ve_{\pm 1/2}$. Clearly, Φ intertwines the actions of $\mathcal{O}(\mathrm{S}_{qr}^2) \rtimes \mathcal{U}_q(\mathrm{su}_2)$ on $M_l \otimes V_{1/2}$ and $\mathcal{O}(\mathrm{SU}_q(2))$. From (42) and the first equality in (43), we conclude that $\Phi(v_{l-1/2, l-1/2}^{l-1/2}) = 0$. Since the highest weight vector $v_{l-1/2, l-1/2}^{l-1/2}$ is cyclic for the representation $\pi_{l-1/2}$, the latter implies that $\Phi(W_{l-1/2}) = \{0\}$. On the other hand, it follows from (17) and (20) that there is a non-zero constant γ_l such that $\Phi(v_{ll}^l \otimes v_{1/2}^{1/2}) = \gamma_l v_{l+1/2, l+1/2}^{l+1/2}$. Therefore, Φ maps $W_{l+1/2}$ into $M_{l+1/2}$, so Φ is a non-trivial intertwiner of the irreducible integrable representations $\pi_{l+1/2}$ on $W_{l+1/2}$ and $\hat{\pi}_{l+1/2}$ on $M_{l+1/2}$. By a standard application of Schur's lemma, $\pi_{l+1/2}$ and $\hat{\pi}_{l+1/2}$ are unitarily equivalent. This proves the assertion for $l+1/2$.

The proof for negative integers is similar. One replaces v_{ll}^l by $v_{-l, -l}^l$ and uses $e_{-1/2} := \|a - q^{2l+1}sc\|^{-1}(a - q^{2l+1}sc)$ and the second equality in (43). $\square$

By Proposition 4.6 and Theorem 3.1, each irreducible integrable $*$ -representation of the cross product algebra $\mathcal{O}(\mathrm{S}_{qr}^2) \rtimes \mathcal{U}_q(\mathrm{su}_2)$ can be realized on $\mathcal{O}(\mathrm{SU}_q(2))$ and $\mathcal{O}(\mathrm{SU}_q(2))$ decomposes into the orthogonal direct sum of all irreducible integrable $*$ -representation of $\mathcal{O}(\mathrm{S}_{qr}^2) \rtimes \mathcal{U}_q(\mathrm{su}_2)$, each with multiplicity one.

5 Representations of the cross product $*$ -algebras: Second approach

5.1 “Decoupling” of the cross product algebras

Let us first suppose that $r \in [0, \infty)$. From the relations $AB = q^{-2}BA$ and $AB^* = q^2B^*A$, it is clear that $\mathcal{S} = \{A^n; n \in \mathbb{N}_0\}$ is a left and right Ore set of the algebra $\mathcal{O}(\mathrm{S}_{qr}^2)$. Moreover the algebra $\mathcal{O}(\mathrm{S}_{qr}^2)$ has no zero divisors. Hence the localization algebra, denoted by $\hat{\mathcal{O}}(\mathrm{S}_{qr}^2)$, of $\mathcal{O}(\mathrm{S}_{qr}^2)$ at $\mathcal{S}$ exists. The $*$ -algebra $\mathcal{O}(\mathrm{S}_{qr}^2)$ is then a $*$ -subalgebra of $\hat{\mathcal{O}}(\mathrm{S}_{qr}^2)$ and all elements $A^n, n \in \mathbb{N}_0$, are invertible in $\hat{\mathcal{O}}(\mathrm{S}_{qr}^2)$. From [9, Theorem 3.4.1], we conclude that $\hat{\mathcal{O}}(\mathrm{S}_{qr}^2)$ is a left $\mathcal{U}_q(\mathrm{su}_2)$ -module $*$ -algebra such that $\mathcal{O}(\mathrm{S}_{qr}^2)$ is a left $\mathcal{U}_q(\mathrm{su}_2)$ -module $*$ -subalgebra. The

left action of the generators E, F, K on A^{-1} is given by

$$E \triangleright A^{-1} = -q^{-5/2} B^* A^{-2}, \quad F \triangleright A^{-1} = q^{1/2} B A^{-2}, \quad K \triangleright A^{-1} = A^{-1}.$$

Hence the left cross product algebra $\hat{\mathcal{O}}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ is a well defined $*$ -algebra containing $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ as a $*$ -subalgebra.

Let $\mathcal{Y}_r$ denote the $*$ -subalgebra of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ generated by

$$X := q^{3/2} \lambda F K^{-1} A + q B = q^{3/2} \lambda A F K^{-1} + q^{-1} B, \quad (44)$$

$$X^* := q^{3/2} \lambda A K^{-1} E + q B^* = q^{3/2} \lambda K^{-1} E A + q^{-1} B^*, \quad (45)$$

$$Y := q K^{-2} A \quad (46)$$

and let $\hat{\mathcal{Y}}_r$ be the subalgebra of $\hat{\mathcal{O}}(S_{qr}^2)$ generated by $\mathcal{Y}_r$ and Y^{-1} . Note that $Y = Y^*$. It is straightforward to check that the elements X, X^*, Y, Y^{-1} commute with the generators A, B, B^* of $\mathcal{O}(S_{qr}^2)$. Hence the algebras $\hat{\mathcal{Y}}_r$ and $\hat{\mathcal{O}}(S_{qr}^2)$ commute inside the cross product algebra $\hat{\mathcal{O}}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$. Moreover, the generators of $\mathcal{Y}_r$ satisfy the commutation relations

$$Y X = q^2 X Y, \quad Y X^* = q^{-2} X^* Y, \quad X^* X - q^2 X X^* = (1 - q^2)(Y^2 + r). \quad (47)$$

We denote by $\hat{\mathcal{U}}_q(\mathfrak{su}_2)$ the Hopf $*$ -subalgebra of $\mathcal{U}_q(\mathfrak{su}_2)$ generated by $e := EK, f := K^{-1}F$ and $k := K^2$. As an algebra, $\hat{\mathcal{U}}_q(\mathfrak{su}_2)$ has generators e, f, k, k^{-1} with defining relations

$$k k^{-1} = k^{-1} k = 1, \quad k e = q^2 e k, \quad k f = q^{-2} f k, \quad e f - f e = \lambda^{-1}(k - k^{-1}). \quad (48)$$

From (44)–(46), we obtain

$$f = q^{-1/2} \lambda^{-1} (X - q B) A^{-1}, \quad e = q^{1/2} \lambda^{-1} (X^* - q^{-1} B^*) Y^{-1}, \quad k = q Y^{-1} A. \quad (49)$$

Hence the two commuting algebras $\hat{\mathcal{Y}}_r$ and $\hat{\mathcal{O}}(S_{qr}^2)$ generate the $*$ -subalgebra $\hat{\mathcal{O}}(S_{qr}^2) \rtimes \hat{\mathcal{U}}_q(\mathfrak{su}_2)$ of $\hat{\mathcal{O}}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$. In this sense, $\hat{\mathcal{Y}}_r$ and $\hat{\mathcal{O}}(S_{qr}^2)$ “decouple” the cross product algebra $\hat{\mathcal{O}}(S_{qr}^2) \rtimes \hat{\mathcal{U}}_q(\mathfrak{su}_2)$. By choosing a PBW-basis of $\hat{\mathcal{O}}(S_{qr}^2)$ and $\hat{\mathcal{Y}}_r$, it is easy to show that the subalgebra generated by $\hat{\mathcal{O}}(S_{qr}^2)$ and $\hat{\mathcal{Y}}_r$ is isomorphic to the tensor product algebra $\hat{\mathcal{O}}(S_{qr}^2) \otimes \hat{\mathcal{Y}}_r$. Therefore we can consider $\hat{\mathcal{O}}(S_{qr}^2) \otimes \hat{\mathcal{Y}}_r$ as a $*$ -subalgebra of $\hat{\mathcal{O}}(S_{qr}^2) \rtimes \hat{\mathcal{U}}_q(\mathfrak{su}_2)$ by identifying $x \otimes y$ with xy , $x \in \hat{\mathcal{O}}(S_{qr}^2)$, $y \in \hat{\mathcal{Y}}_r$. Similarly, $\mathcal{O}(S_{qr}^2) \otimes \mathcal{Y}_r$ becomes a $*$ -subalgebra of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$.

There is an alternative way to define the $*$ -algebra $\hat{\mathcal{O}}(S_{qr}^2) \rtimes \hat{\mathcal{U}}_q(\mathfrak{su}_2)$ by taking the two sets $A = A^*, A^{-1}, B, B^*$ and $X, X^*, Y = Y^*, Y^{-1}$ of pairwise commuting generators with defining relations (11), (47) and the obvious relations

$$AA^{-1} = A^{-1}A = 1, \quad YY^{-1} = Y^{-1}Y = 1. \quad (50)$$

Indeed, if we define e, f, k by (49), then the relations (48) of $\mathcal{U}_q(\mathfrak{su}_2)$ and the cross relations of $\hat{\mathcal{O}}(S_{qr}^2) \rtimes \hat{\mathcal{U}}_q(\mathfrak{su}_2)$ can be derived from this set of defining relations.

The larger cross product algebra $\hat{\mathcal{O}}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ can be redefined in a similar manner if we replace the generator Y by K . That is, $\hat{\mathcal{O}}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ is the $*$ -algebra with the generators $A = A^*, A^{-1}, B, B^*, X, X^*, K = K^*, K^{-1}$ satisfying the defining relations (11), (50),

$$KA = AK, \quad BK = qKB, \quad KB^* = qB^*K, \quad X^*X - q^2XX^* = (1 - q^2)(q^2K^{-4}A^2 + r), \quad (51)$$

and A, B, B^* commute with X and X^* . The generators F and E are then given by

$$F = q^{-3/2}\lambda^{-1}(X - qB)KA^{-1}, \quad E = q^{-3/2}\lambda^{-1}A^{-1}K(X^* - qB^*). \quad (52)$$

The preceding considerations and facts carry over almost verbatim to the case $r = \infty$. The only difference is that in the case $r = \infty$ one has to set $r = 1$ in the third equations of (47) and (51).

5.2 Operator representations of the $*$ -algebra $\mathcal{Y}_r$

For the study of representations of the $*$ -algebra $\mathcal{Y}_r$, we need two auxiliary lemmas. The first one restates the *Wold decomposition* of an isometry (see [18, Theorem 1.1]), while the second is Lemma 4.2(ii) in [15].

Lemma 5.1 *Each isometry v on a Hilbert space $\mathcal{K}$ is up to unitary equivalence of the following form: There exist Hilbert subspaces $\mathcal{K}^u$ and $\mathcal{K}_0^s$ of $\mathcal{K}$ and a unitary operator v_u on $\mathcal{K}^u$ such that $v = v_u \oplus v_s$ on $\mathcal{K} = \mathcal{K}^u \oplus \mathcal{K}^s$ and v_s acts on $\mathcal{K}^s = \bigoplus_{n=0}^{\infty} \mathcal{K}_n^s$ by $v_s \zeta_n = \zeta_{n+1}$, where each $\mathcal{K}_n^s$ is $\mathcal{K}_0^s$ and $\zeta_n \in \mathcal{K}_n^s$. Moreover, $\mathcal{K}^u = \bigcap_{n=0}^{\infty} v^n \mathcal{K}$.*

Lemma 5.2 *Let v be the operator v_s on $\mathcal{K}^s = \bigoplus_{n=0}^{\infty} \mathcal{K}_n^s$, $\mathcal{K}_n^s = \mathcal{K}_0^s$, from Lemma 5.1 and let Y be a self-adjoint operator on $\mathcal{K}^s$ such that $q^2 v Y \subseteq Y v$. Then there is a self-adjoint operator Y_0 on the Hilbert space $\mathcal{K}_0^s$ such that $Y \zeta_n = q^{2n} Y_0 \zeta_n$ and $v \zeta_n = \zeta_{n+1}$ for $\zeta_n \in \mathcal{K}_n^s$ and $n \in \mathbb{N}_0$.*

It suffices to treat the case $r \in [0, \infty)$ because the algebra $\mathcal{Y}_\infty$ is isomorphic to $\mathcal{Y}_1$. Suppose that we have a representation of relations (47) by closed operators X , X^* and a self-adjoint operator Y acting on a Hilbert space $\mathcal{K}$. We assume that Y has trivial kernel.

Let $X = v|X|$ be the polar decomposition of the operator X . Since $q \in (0, 1)$, $r \geq 0$ and $\ker Y = \{0\}$, we have $\ker X = \{0\}$ by the third equation of (47). Hence v is an isometry on $\mathcal{K}$, that is, $v^*v = I$.

By (47), $YX^*X = X^*XY$. We assume that the self-adjoint operators Y and X^*X strongly commute. Then Y and $|X| = (X^*X)^{1/2}$ also strongly commute. Again by (47), $Yv|X| = YX = q^2XY = q^2v|X|Y = q^2vY|X|$. Since $\ker |X| = \ker X = \{0\}$, we conclude that

$$q^2vY = Yv. \quad (53)$$

Since $X^* = |X|v^*$, the third equation of (47) rewrites as

$$|X|^2 = q^2v|X|^2v^* + (1-q^2)(Y^2 + r). \quad (54)$$

Multiplying by v gives $|X|^2v = v(q^2|X|^2 + (1-q^2)(q^4Y^2 + r))$ since $v^*v = I$ and, by (53), $Y^2v = q^4vY^2$. Proceeding by induction, we derive

$$|X|^2v^n = v^n(q^{2n}|X|^2 + (1-q^{2n})(q^{2n+2}Y^2 + r)). \quad (55)$$

Now we use the Wold decomposition $v = v_u \oplus v_s$ on $\mathcal{K} = \mathcal{K}^u \oplus \mathcal{K}^s$ of the isometry v by Lemma 5.1. Since $\mathcal{K}^u = \cap_{n=0}^\infty v^n \mathcal{K}$ and $Yv = q^2vY$, the Hilbert subspace $\mathcal{K}^u$ reduces the self-adjoint operator Y . From (55), it follows that $|X|^2$ leaves a dense subspace of the space $\mathcal{K}^u$ invariant. We assume that $\mathcal{K}^u$ reduces $|X|^2$. Hence we can consider all operators occurring in relation (55) on the subspace $\mathcal{K}^u$. For the unitary part v_u of v , we have $v_u v_u^* = I$ on $\mathcal{K}^u$. Multiplying (55) by $(v_u^*)^n$ and using again Equation (53), we derive on $\mathcal{K}^u$ the relation

$$0 \leq q^{2n}v_u^n |X|^2 (v_u^*)^n = |X|^2 - (1-q^{2n})r + q^2Y^2 - q^{-2n+2}Y^2$$

for all $n \in \mathbb{N}$. Letting $n \rightarrow \infty$ and remembering that $\ker Y = \{0\}$ and $q \in (0, 1)$, we conclude that the latter is only possible when $\mathcal{K}^u = \{0\}$. That is, the isometry v is (unitarily equivalent to) the unilateral shift operator v_s .

Since $v = v_s$, Lemma 5.2 applies to the relation $q^2vY = Yv$. Hence there exists a self-adjoint operator Y_0 on $\mathcal{K}_0^s$ such that $Y\zeta_n = q^{2n}Y_0\zeta_n$ on $\mathcal{K}^s = \bigoplus_{n=0}^\infty \mathcal{K}_n^s$,

$\mathcal{K}_n^s = \mathcal{K}_0^s$. Since $v^*\zeta_0 = 0$, (54) yields $|X|^2\zeta_0 = (1 - q^2)(Y_0^2 + r)\zeta_0$. Using this equation and (54), we compute

$$\begin{aligned} |X|^2\zeta_n &= |X|^2v^n\zeta_0 = v^n(q^{2n}|X|^2 + (1 - q^{2n})(q^{2n+2}Y^2 + r))\zeta_0 \\ &= v^n(q^{2n}(1 - q^2)(Y_0^2 + r)\zeta_0 + (1 - q^{2n})(q^{2n+2}Y_0^2 + r)\zeta_0) \\ &= \lambda_{n+1}^2(q^{2n}Y_0^2 + r)\zeta_n \end{aligned} \quad (56)$$

for $\zeta_n \in \mathcal{K}_n^s$. We assumed above that the pointwise commuting self-adjoint operator $|X|^2$ and Y strongly commute. Hence their reduced parts on each subspace $\mathcal{K}_n^s$ strongly commute. Therefore, (56) implies $|X|\zeta_n = \lambda_{n+1}(q^{2n}Y_0^2 + r)^{1/2}\zeta_n$. Recall that $X = v|X|$ and $X^* = |X|v^*$. Renaming $\mathcal{K}_n^s$ by $\mathcal{K}_n$ and summarizing the preceding, we obtain the following form of the operators X , X^* and Y :

$$\begin{aligned} X\zeta_n &= \lambda_{n+1}(q^{2n}Y_0^2 + r)^{1/2}\zeta_{n+1}, \quad X^*\zeta_n = \lambda_n(q^{2n-2}Y_0^2 + r)^{1/2}\zeta_{n-1}, \\ Y\zeta_n &= q^{2n}Y_0\zeta_n \quad \text{on } \mathcal{K} = \bigoplus_{n=0}^{\infty} \mathcal{K}_n, \quad \mathcal{K}_n = \mathcal{K}_0, \end{aligned}$$

where Y_0 is self-adjoint operator with trivial kernel on the Hilbert space $\mathcal{K}_0$. Conversely, it is easy to check that these operators X , X^* , Y satisfy the relations (47), so they define indeed a $*$ -representation of the $*$ -algebra $\mathcal{Y}_r$. The representation is irreducible if and only if $\mathcal{K}_0 = \mathbb{C}$. In this case, Y_0 is a non-zero real number.

5.3 Representations of the cross product $*$ -algebras

First let us review the representations of the $*$ -algebra $\mathcal{O}(S_{qr}^2)$ from [13]. Recall the definition of $\lambda_{\pm}$ from Section 3. For $r < \infty$, we set

$$c_{\pm}(n) := (r + \lambda_{\pm}q^{2n} - (\lambda_{\pm}q^{2n})^2)^{1/2}.$$

Let $\mathcal{H}^0$, $\mathcal{H}_0^+$ and $\mathcal{H}_0^-$ be Hilbert spaces and let u be a unitary operator on $\mathcal{H}^0$. Let $\mathcal{H}^{\pm} = \bigoplus_{n=0}^{\infty} \mathcal{H}_n^{\pm}$, where $\mathcal{H}_n^{\pm} = \mathcal{H}_0^{\pm}$, and let $\mathcal{H} = \mathcal{H}^0 \oplus \mathcal{H}^+ \oplus \mathcal{H}^-$ for $r \in (0, \infty]$ and $\mathcal{H} = \mathcal{H}^0 \oplus \mathcal{H}^+$ for $r = 0$. The generators of $\mathcal{O}(S_{qr}^2)$ act on the Hilbert space $\mathcal{H}$ by the following formulas:

$$\begin{aligned} r \in [0, \infty) : \quad & A = 0, \quad B = r^{1/2}u, \quad B^* = r^{1/2}u^* \quad \text{on } \mathcal{H}^0, \\ & A\eta_n = \lambda_{\pm}q^{2n}\eta_n, \quad B\eta_n = c_{\pm}(n)\eta_{n-1}, \quad B^*\eta_n = c_{\pm}(n+1)\eta_{n+1} \quad \text{on } \mathcal{H}^{\pm}. \\ r = \infty : \quad & A = 0, \quad B = u, \quad B^* = u^* \quad \text{on } \mathcal{H}_0, \\ & A\eta_n = \pm q^{2n}\eta_n, \quad B\eta_n = (1 - q^{4n})^{1/2}\eta_{n-1}, \quad B^*\eta_n = (1 - q^{4(n+1)})^{1/2}\eta_{n+1} \quad \text{on } \mathcal{H}^{\pm}. \end{aligned}$$

Recall that, for $r = 0$, there is no Hilbert space $\mathcal{H}^-$. From [13, Proposition 4], it follows that, up to unitary equivalence, each $*$ -representation of $\mathcal{O}(S_{qr}^2)$ is of the above form. Note that all operators are bounded and $\mathcal{H}^0 = \ker A$, $A > 0$ on $\mathcal{H}^+$ and $A < 0$ on $\mathcal{H}^-$.

Next we show that, for each $*$ -representation of the algebras $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ and $\mathcal{O}(S_{qr}^2) \rtimes \hat{\mathcal{U}}_q(\mathfrak{su}_2)$, the space $\mathcal{H}^0$ is $\{0\}$, so the operator A is invertible. We carry out the reasoning for $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$. Assuming that the commuting self-adjoint operators A and K strongly commute, it follows that K leaves $\mathcal{H}^0 = \ker A$ invariant. By (11) (resp. (12)), $B\mathcal{H}^0 \subseteq \mathcal{H}^0$. Let $\xi \in \mathcal{H}^0$. Using the relation $FA = AF - q^{-3/2}BK$, we see that

$$q^{-3}\|BK\xi\|^2 = \|AF\xi\|^2 = \langle q^{-3/2}BK\xi, AF\xi \rangle = \langle q^{-3/2}ABK\xi, F\xi \rangle = 0.$$

That is, $BK\xi = 0$ and $AF\xi = 0$, so that $F\xi \in \mathcal{H}^0$. For $r \in (0, \infty]$, BK is invertible on $\mathcal{H}^0$ and hence $\xi = 0$. For $r = 0$, we have $B^*\xi = B^*F\xi = 0$. From the cross relation $FB^* = q^{-1}B^*F + q^{-1/2}(1+q^2)AK - q^{-1/2}K$, we get $K\xi = 0$ and so $\xi = 0$. Thus, $\mathcal{H}^0 = \{0\}$.

Consider now a $*$ -representation of the $*$ -algebra $\hat{\mathcal{O}}(S_{qr}^2) \rtimes \hat{\mathcal{U}}_q(\mathfrak{su}_2)$. Its restriction to $\mathcal{O}(S_{qr}^2)$ is a $*$ -representation of the form described above with $\mathcal{H}^0 = \{0\}$. All operators of the $*$ -subalgebra $\mathcal{Y}_r$ commute with A and B . Let us assume that the spectral projections of the self-adjoint operator A commute also with all operators of $\mathcal{Y}_r$. (Note that, for a $*$ -representation by bounded operators on a Hilbert space, the latter fact follows. For unbounded $*$ -representation, it does not and we restrict ourselves to the class of well behaved representations which satisfy this assumption.) Since $\mathcal{H}_n^\pm$ is the eigenspace of A with eigenvalue $\lambda_\pm q^{2n}$ and these eigenvalues are pairwise distinct, the operators of $\mathcal{Y}_r$ leave $\mathcal{H}_n^\pm$ invariant. That is, we have a $*$ -representation of $\mathcal{Y}_r$ on $\mathcal{H}_n^\pm$. But $\mathcal{Y}_r$ commutes also with B . Since B is a weighted shift operators with weights $c_\pm(n) \neq 0$ for $n \in \mathbb{N}$, it follows that the representations of $\mathcal{Y}_r$ on $\mathcal{H}_n^\pm = \mathcal{H}_0^\pm$ are the same for all $n \in \mathbb{N}_0$. Using the structure of the representation of the $*$ -algebra $\mathcal{Y}_r$ on the Hilbert space $\mathcal{K} := \mathcal{H}_0^\pm$ derived in Subsection 5.2 and inserting the formulas for X, X^*, Y and A, B, B^* into (49), one obtains the action of the generators e, f, k, k^{-1} of $\hat{\mathcal{U}}_q(\mathfrak{su}_2)$. We do not list these formulas, but we will do so below for the generators of $\mathcal{U}_q(\mathfrak{su}_2)$.

Let us turn to a $*$ -representation of the larger $*$ -algebra $\hat{\mathcal{O}}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$. Then, in addition to the considerations of the preceding paragraph, we have to deal with the generator K . We assumed above that K and A are strongly commuting self-adjoint operators. Therefore, K commutes with the spectral projections of A . Hence each space $\mathcal{H}_n^\pm$ is reducing for K . The relation $BK = qKB$ implies that

there is an invertible self-adjoint operator K_0 on $\mathcal{H}_0^\pm$ such that $K\eta_n = q^n K_0 \eta_n$ for $\eta_n \in \mathcal{H}_n^\pm$. Recall that we have $XK = qKX$ and $YK = KY$ in the algebra $\hat{\mathcal{O}}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$. Inserting for X and Y the corresponding operators on $\mathcal{K} \equiv \mathcal{H}_0^\pm = \bigoplus_{m=0}^\infty \mathcal{K}_m$ from Subsection 5.2, we conclude that there exists an invertible self-adjoint operator H on $\mathcal{K}_0$ such that $K_0 \zeta_m = q^{-m} H \zeta_m$ for $\zeta_m \in \mathcal{K}_m$. Further, since $Y = qK^{-2}A$, we have $Y\zeta_0 = Y_0 \zeta_0 = qH^{-2} \lambda_\pm \zeta_0$ for $\zeta_0 \in \mathcal{K}_0$. Inserting the preceding facts and the results from Subsection 5.2 into (52) and renaming $\mathcal{K}_0$ by $\mathcal{G}$, we obtain the following $*$ -representations (satisfying the assumptions made above) of the cross product $*$ -algebra $\hat{\mathcal{O}}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ for $r \in (0, \infty)$:

$$\begin{aligned} (I)_{\pm, H} : \quad & A\eta_{nm} = \lambda_\pm q^{2n} \eta_{nm}, \quad B\eta_{nm} = c_\pm(n) \eta_{n-1, m}, \quad B^* \eta_{nm} = c_\pm(n+1) \eta_{n+1, m}, \\ & E\eta_{nm} = q^{-1/2} \lambda^{-1} [q^{-n} \lambda_m (\lambda_\pm^{-2} q^{-2m} r + H^{-4})^{1/2} H \eta_{n, m-1} \\ & \quad - q^{-m} (\lambda_\pm^{-2} q^{-2n-2} r + \lambda_\pm^{-1} - q^{2n+2})^{1/2} H \eta_{n+1, m}], \\ & F\eta_{nm} = q^{-1/2} \lambda^{-1} [q^{-n} \lambda_{m+1} (\lambda_\pm^{-2} q^{-2m-2} r + H^{-4})^{1/2} H \eta_{n, m+1} \\ & \quad - q^{-m} (\lambda_\pm^{-2} q^{-2n} r + \lambda_\pm^{-1} - q^{2n})^{1/2} H \eta_{n-1, m}], \\ & K\eta_{nm} = q^{n-m} H \eta_{nm}, \end{aligned}$$

where H is an invertible self-adjoint operator on a Hilbert space $\mathcal{G}$. The domain is the direct sum $\mathcal{H} = \bigoplus_{n, m=0}^\infty \mathcal{H}_{nm}$, where $\mathcal{H}_{nm} = \mathcal{G}$. In the case $r = 0$, there is only the representation $(I)_{+, H}$. The case $r = \infty$ has already been treated in [15].

6 Algebras of functions and representations of the cross product algebras

6.1 Algebras of functions and the invariant state on S_{qr}^2

Let $\mathcal{F}(\sigma(A))$ be the $*$ -algebra of all complex Borel functions on the set $\sigma(A) := \{q^{2n} \lambda_+, q^{2n} \lambda_-; n \in \mathbb{N}_0\}$, let $\mathcal{F}_b(\sigma(A))$ be the $*$ -algebra of all bounded complex Borel functions on $\sigma(A)$, and let $\mathcal{F}^\infty(\sigma(A))$ be the set of all $f \in \mathcal{F}(\sigma(A))$ for which there exist an $\varepsilon > 0$ and a function $\tilde{f} \in C^\infty(-\varepsilon, \varepsilon)$ such that $f = \tilde{f}$ on $\sigma(A) \cap (-\varepsilon, \varepsilon)$. For $f \in \mathcal{F}^\infty(\sigma(A))$, we can assign unambiguously the value $f(0)$ at 0 by taking the limit $f(0) = \lim_{t \rightarrow 0} f(t)$. In order to be in accordance with our previous notation, we write A for the function $\text{id}(t) = t$ and also for the argument of functions $f \in \mathcal{F}(\sigma(A))$. We denote by $\mathcal{F}(S_{qr}^2)$, $r \in [0, \infty]$, the unital $*$ -algebra generated by the $*$ -algebra $\mathcal{F}(\sigma(A))$ and two generators B, B^* with defining

relations

$$Bf(A) = f(q^2 A)B, \quad f(A)B^* = B^*f(q^2 A), \quad f \in \mathcal{F}(\sigma(A)); \quad (57)$$

$$B^*B = A - A^2 + r, \quad BB^* = q^2 A - q^4 A^2 + r, \quad \text{for } r \in [0, \infty), \quad (58)$$

$$B^*B = -A^2 + 1, \quad BB^* = -q^4 A^2 + 1 \quad \text{for } r = \infty. \quad (59)$$

For $k \in \mathbb{Z}$, we set $B^{\#k} = B^k$ if $k \geq 0$ and $B^{\#k} = B^{*k}$ if $k < 0$. As a vector space, $\mathcal{F}(S_{qr}^2)$ is spanned by $\{B^n f_1(A), f_2(A)B^{*k}; f_1, f_2 \in \mathcal{F}(\sigma(A)), n, k \in \mathbb{N}_0\} = \{f(A)B^{\#k}; f \in \mathcal{F}(\sigma(A)), k \in \mathbb{Z}\}$. We denote by $\mathcal{F}_b(S_{qr}^2)$ and $\mathcal{F}^\infty(S_{qr}^2)$ the $*$ -subalgebras of $\mathcal{F}(S_{qr}^2)$ generated by $\mathcal{F}_b(\sigma(A))$ and $\mathcal{F}^\infty(\sigma(A))$, respectively, and B and B^* . We introduce a left action $\triangleright$ of the Hopf algebra $\mathcal{U}_q(\mathfrak{su}_2)$ on $\mathcal{F}^\infty(S_{qr}^2)$ by setting

$$\begin{aligned} E \triangleright p(B)f(A) &= q^{1/2} \left[\frac{p(q^{-1}B) - p(qB)}{(q^{-1} - q)B} - (1+q^2) \frac{p(q^{-3}B) - p(qB)}{(q^{-3} - q)B} A \right] f(A) \\ &\quad + q^{-1/2} p(qB)B^* \frac{f(A) - f(q^2 A)}{(1 - q^2)A}, \end{aligned}$$

$$E \triangleright f(A)p(B^*) = q^{-1/2} \frac{f(q^{-2}A) - f(A)}{(q^{-2} - 1)A} B^* p(qB^*),$$

$$F \triangleright p(B)f(A) = -q^{-3/2} Bp(qB) \frac{f(q^{-2}A) - f(A)}{(q^{-2} - 1)A},$$

$$\begin{aligned} F \triangleright f(A)p(B^*) &= -q^{-3/2} \frac{f(A) - f(q^2 A)}{(1 - q^2)A} Bp(qB^*) \\ &\quad - q^{-1/2} f(A) \left[\frac{p(q^{-1}B^*) - p(qB^*)}{(q^{-1} - q)B^*} - (1+q^2)A \frac{p(q^{-3}B^*) - p(qB^*)}{(q^{-3} - q)B^*} \right], \end{aligned}$$

$$K \triangleright p(B)f(A) = p(q^{-1}B)f(A), \quad K \triangleright f(A)p(B^*) = f(A)p(qB^*),$$

where f is a function in $\mathcal{F}^\infty(\sigma(A))$ and p is a polynomial. It can be shown that these formulas define indeed an action of the Hopf $*$ -algebra $\mathcal{U}_q(\mathfrak{su}_2)$ such that $\mathcal{F}^\infty(S_{qr}^2)$ is a left $\mathcal{U}_q(\mathfrak{su}_2)$ -module $*$ -algebra. We omit the details of these lengthy, but straightforward computations.

From the defining relations, it is clear that the coordinate $*$ -algebra $\mathcal{O}(S_{qr}^2)$ of the quantum sphere is a $*$ -subalgebra of $\mathcal{F}^\infty(S_{qr}^2)$. On B, A, B^* considered as

elements of $\mathcal{F}^\infty(S_{qr}^2)$, the preceding formulas coincide with corresponding formulas for the actions on the generators B, A, B^* of $\mathcal{O}(S_{qr}^2)$. Hence $\mathcal{O}(S_{qr}^2)$ is a left $\mathcal{U}_q(\mathfrak{su}_2)$ -module $*$ -subalgebra of $\mathcal{F}^\infty(S_{qr}^2)$.

Now we turn to the construction of a $\mathcal{U}_q(\mathfrak{su}_2)$ -invariant linear functional h . For $f \in \mathcal{F}_b(\sigma(A))$, we put

$$h_0(f(A)) := \gamma_+ \sum_{n=0}^{\infty} f(\lambda_+ q^{2n}) q^{2n} + \gamma_- \sum_{n=0}^{\infty} f(\lambda_- q^{2n}) q^{2n},$$

where $\gamma_+ := (1 - q^2)\lambda_+(\lambda_+ - \lambda_-)^{-1}$ and $\gamma_- := (1 - q^2)\lambda_-(\lambda_- - \lambda_+)^{-1}$. Note that $\gamma_{\pm} = (1 - q^2)(1/2 \pm (r + 4)^{-1/2})$. (When $r = 0$, the above equation simplifies to $h_0(f(A)) = (1 - q^2) \sum_{n=0}^{\infty} f(q^{2n}) q^{2n}$ since in this case $\gamma_+ = \lambda_+ = 1$ and $\gamma_- = 0$.) Define a functional h on $\mathcal{F}_b(S_{qr}^2)$ by

$$h(p(B)f(A)) = p(0)h_0(f(A)), \quad h(f(A)p(B^*)) = p(0)h_0(f(A)). \quad (60)$$

For $g \in \mathcal{F}_b(\sigma(A))$, we have

$$h(g(A)) = q^2 h(g(q^2 A)) + \gamma_+ g(\lambda_+) + \gamma_- g(\lambda_-). \quad (61)$$

First we show that h is a faithful state on the $*$ -algebra $\mathcal{F}_b(S_{qr}^2)$. We restrict ourselves to the case $r \in (0, \infty]$. Let $x = \sum_k (B^k f_k(A) + g_k(A) B^{*k}) \in \mathcal{F}_b(S_{qr}^2)$ with $f_k, g_k \in \mathcal{F}_b(\sigma(A))$. From (57)–(59), we obtain $B^{*k} B^k = \prod_{i=0}^{k-1} p_r(q^{-2i} A)$ and $B^k B^{*k} = \prod_{i=1}^k p_r(q^{2i} A)$, where $p_r(A) := (\lambda_+ - A)(A - \lambda_-)$. By (60), we have

$$\begin{aligned} h(x^* x) &= \sum_k h(B^k \overline{g_k} g_k B^{*k} + \overline{f_k} B^{*k} B^k f_k) \\ &= \sum_k h_0(|g_k(q^{2k} A)|^2 \prod_{i=1}^k p_r(q^{2i} A)) + h_0(|f_k(A)|^2 \prod_{i=0}^{k-1} p_r(q^{-2i} A)) \end{aligned} \quad (62)$$

Note that $p_r(\lambda_{\pm}) = 0$ and $p_r(q^{2j} \lambda_{\pm}) > 0$ for $j \in \mathbb{N}$. Hence $h(x^* x) \geq 0$ by (62). Assume that $h(x^* x) = 0$. From (62) and the definition of h_0 , it follows that $g_k(q^{2n} \lambda_{\pm}) = 0$ for $n \in \mathbb{N}_0$ and $f_k(q^{2n} \lambda_{\pm}) = 0$ for $n \geq k$. The latter implies $g_k(A) = 0$ and $B^k f_k(A) = 0$, so $x = 0$. That is, h is a faithful state.

Now we prove that h is $\mathcal{U}_q(\mathfrak{su}_2)$ -invariant on the $*$ -subalgebra $\mathcal{F}^\infty(S_{qr}^2)$, that is, $h(y \triangleright x) = \varepsilon(y)h(x)$ for $y \in \mathcal{U}_q(\mathfrak{su}_2)$ and $x \in \mathcal{F}^\infty(S_{qr}^2)$. Clearly, it suffices to verify the latter condition for the generators $y = E, F, K$. For $y = K$, the assertion is obvious. Since the functional h is hermitian and $\mathcal{F}^\infty(S_{qr}^2)$ is a $\mathcal{U}_q(\mathfrak{su}_2)$ -module $*$ -algebra, it is sufficient to check the invariance with respect to E . By

the definitions of the action of E and the functional h , it is enough to show that $h(E \triangleright Bf(A)) = 0$ for all $f(A) \in \mathcal{F}^\infty(S_{qr}^2)$. We carry out the proof for $r \in (0, \infty)$. Since $f \in \mathcal{F}^\infty(S_{qr}^2)$, the function $g(A) := f(A) - Af(A) + r(f(A) - f(0))/A$ is bounded on $\sigma(A)$. Hence $g \in \mathcal{F}_b(\sigma(A))$ and (61) applies to this function. Observe that $1 - \lambda_\pm + r\lambda_\pm^{-1} = 0$ and $\gamma_+\lambda_+^{-1} + \gamma_-\lambda_-^{-1} = 0$ by the definitions of these constants. Using these facts and the equation $BB^* = q^2A - q^4A^2 + r$, we compute

$$\begin{aligned} q^{-1/2}E \triangleright Bf(A) &= f(A) - (1 + q^2)Af(A) + BB^* \frac{f(A) - f(q^2A)}{(1 - q^2)A} \\ &= (1 - q^2)^{-1}(g(A) - g(q^2A)q^2) \end{aligned}$$

and hence

$$\begin{aligned} q^{-1/2}(1 - q^2)h(E \triangleright Bf(A)) &= \gamma_+g(\lambda_+) + \gamma_-g(\lambda_-) = \\ \gamma_+f(\lambda_+)(1 - \lambda_+ + r\lambda_+^{-1}) &+ \gamma_-f(\lambda_-)(1 - \lambda_- + r\lambda_-^{-1}) - f(0)(\gamma_+\lambda_+^{-1} + \gamma_-\lambda_-^{-1}) = 0. \end{aligned}$$

Thus, h is a $\mathcal{U}_q(\mathfrak{su}_2)$ -invariant state on $\mathcal{F}^\infty(S_{qr}^2)$.

For the coordinate algebra $\mathcal{O}(S_{qr}^2)$, the preceding description of the invariant functional was obtained in [10]. However for our consideration below it is crucial to have the invariant state on the larger $*$ -algebra $\mathcal{F}^\infty(S_{qr}^2)$.

The preceding proof shows that, with the action of E defined by the above formula, $h(E \triangleright Bf(A)) = 0$ for $f \in \mathcal{F}_b(\sigma(A))$ if $f(0) := \lim_{t \rightarrow 0} f(t)$ exists and the function $(f(A) - f(0))/A$ is bounded on $\sigma(A)$. For instance, $h(E \triangleright BA\chi_+(A)) = 0$, where χ_+ is the characteristic function of $[0, \infty)$. But we get $h(E \triangleright B\chi_+(A)) = q^{1/2}(1 - q^2)^{-1}\gamma_+\lambda_- \neq 0$ for $r \in (0, \infty]$, so the $\mathcal{U}_q(\mathfrak{su}_2)$ -invariance of the functional h does not hold on the larger $*$ -algebra $\mathcal{F}_b(S_{qr}^2)$.

Now we develop a second operator-theoretic approach to the $\mathcal{U}_q(\mathfrak{su}_2)$ -module structure of $\mathcal{F}^\infty(S_{qr}^2)$ and to the invariant state h . Suppose that π is a $*$ -representation of the $*$ -algebra $\mathcal{F}_b(S_{qr}^2)$ on a Hilbert space $\mathcal{H}$ such that $\ker \pi(A) = \{0\}$. Then all operators $\pi(x)$, $x \in \mathcal{F}_b(S_{qr}^2)$, are bounded and leave the dense domain $\mathcal{D} := \cap_{n=1}^\infty \mathcal{D}(\pi(A)^{-n})$ of $\mathcal{H}$ invariant. For notational simplicity, we write x for the operators $\pi(x)$ and $\pi(x)|\mathcal{D}$. From the form of the representations of $\mathcal{O}(S_{qr}^2)$ described in Subsection 5.3, it is clear that $\text{sign} A := A|A|^{-1}$ commutes with all representation operators. For $T \in \mathcal{L}^+(\mathcal{D})$, we define

$$\begin{aligned} E \triangleright T &= -q^{1/2}\lambda^{-1}A^{-1}[B^*, |A|^{1/2}T|A|^{-1/2}] \\ &= -q^{-1/2}\lambda^{-1}\text{sign} A |A|^{-1/2}[B^*, T]|A|^{-1/2}, \end{aligned} \tag{63}$$

$$\begin{aligned} F \triangleright T &= -q^{-5/2} \lambda^{-1} A^{-1} [B, |A|^{1/2} T |A|^{-1/2}] \\ &= -q^{-3/2} \lambda^{-1} \text{sign} A |A|^{-1/2} [B, T] |A|^{-1/2}, \end{aligned} \quad (64)$$

$$K \triangleright T = |A|^{1/2} T |A|^{-1/2}, \quad K^{-1} \triangleright T = |A|^{-1/2} T |A|^{1/2}. \quad (65)$$

Then one verifies that the $*$ -algebra $\mathcal{L}^+(\mathcal{D})$ is a $\mathcal{U}_q(\mathfrak{su}_2)$ -module $*$ -algebra with the actions of the generators E, F, K, K^{-1} given by these formulas and $\mathcal{F}^\infty(S_{qr}^2)$ is a $\mathcal{U}_q(\mathfrak{su}_2)$ -module $*$ -subalgebra of $\mathcal{L}^+(\mathcal{D})$. Setting $T = p(B)f(A)$ or $T = f(A)p(B^*)$, one easily obtains the formulas for the actions of E, F, K, K^{-1} on $\mathcal{F}^\infty(S_{qr}^2)$ listed above.

In order to define the invariant state h , we specialize the representation π . For $r \in (0, \infty]$, let π be the direct sum representation on $\mathcal{H} := \mathcal{H}^+ \oplus \mathcal{H}^-$, where the representations on $\mathcal{H}^\pm$ are given in Subsection 5.3 with $\mathcal{H}_0^\pm = \mathbb{C}$. For $r = 0$, we take the representation on $\mathcal{H} := \mathcal{H}^+$ with $\mathcal{H}_0^+ = \mathbb{C}$. Since the operator A has an orthonormal basis of eigenvectors with eigenvalues $q^{2n} \lambda_\pm$ and each eigenvalue has multiplicity one, A is of trace class and so is $|A|x$ for all $x \in \mathcal{F}_b(S_{qr}^2)$. Obviously, $\text{Tr}_{\mathcal{H}} |A| = (1 - q^2)^{-1} (\lambda_+ - \lambda_-)$. Therefore,

$$h(x) = (1 - q^2)(\lambda_+ - \lambda_-)^{-1} \text{Tr}_{\mathcal{H}} |A|x, \quad x \in \mathcal{F}_b(S_{qr}^2), \quad (66)$$

defines a state on the $*$ -algebra $\mathcal{F}_b(S_{qr}^2)$.

Next we show that h is $\mathcal{U}_q(\mathfrak{su}_2)$ -invariant on $\mathcal{F}^\infty(S_{qr}^2)$. We carry out this verification in the case $r \in (0, \infty]$ for the generator E and for an element $x = B^{\#n} f(A)$ of $\mathcal{F}^\infty(S_{qr}^2)$, where $n \in \mathbb{Z}$ and $f \in \mathcal{F}^\infty(\sigma(A))$. The other cases are treated in a similar manner. Since $f \in \mathcal{F}^\infty(\sigma(A))$, there is a bounded function g on $\sigma(A)$ such that $f(A) - f(0) = Ag(A)$. Write $x = x_1 + x_2$, where $x_1 = B^{\#n} g(A)A$ and $x_2 = f(0)B^{\#n}$. By (63) and (66), we have

$$h(E \triangleright y) = \text{const} \text{Tr}_{\mathcal{H}} |A| A^{-1} [B^*, y] = \text{const} (\text{Tr}_{\mathcal{H}^+} [B^*, y] - \text{Tr}_{\mathcal{H}^-} [B^*, y]) \quad (67)$$

for $y \in \mathcal{F}^\infty(S_{qr}^2)$. Since $g(A)$ and $B^{\#n}$ are bounded operators and A is of trace class, x_1 is of trace class. Therefore, $\text{Tr}_{\mathcal{H}^\pm} [B^*, x_1] = 0$, so that $h(E \triangleright x_1) = 0$ by (67). Since $[B^*, B^{\#n}] \in A \cdot \mathcal{O}(S_{qr}^2)$ by Lemma 6.2 below, $[B^*, B^{\#n}]$ is of trace class. If $n \neq 1$, then $\text{Tr}_{\mathcal{H}} |A| A^{-1} [B^*, B^{\#n}] = 0$ because in this case $|A| A^{-1} [B^*, B^{\#n}]$ acts on $\mathcal{H}$ as a weighted shift operator. Hence $h(E \triangleright x_2) = 0$. Suppose that $n = 1$. Since $[B^*, B] = (1 - q^2)(A - (1 + q^2)A^2)$ and

$$(1 - q^2) \text{Tr}_{\mathcal{H}^\pm} (A - (1 + q^2)A^2) = \lambda_\pm - \lambda_\pm^2 = \lambda_+ \lambda_-,$$

it follows from (67) that $h(E \triangleright x_2) = 0$. Thus, $h(E \triangleright x) = 0$. This completes the proof of the $\mathcal{U}_q(\mathfrak{su}_2)$ -invariance of h on $\mathcal{F}^\infty(S_{qr}^2)$.

In both approaches, we have proved the following theorem.

Theorem 6.1 *With the foregoing definitions, $\mathcal{F}^\infty(S_{qr}^2)$ is a left $\mathcal{U}_q(\mathfrak{su}_2)$ -module $*$ -algebra which contains $\mathcal{O}(S_{qr}^2)$ as a $\mathcal{U}_q(\mathfrak{su}_2)$ -module $*$ -subalgebra. The functional h is a faithful $\mathcal{U}_q(\mathfrak{su}_2)$ -invariant state on the $\mathcal{U}_q(\mathfrak{su}_2)$ -module $*$ -algebra $\mathcal{F}^\infty(S_{qr}^2)$.*

In the next subsection, we shall need the following lemma. It is the algebraic version of the operator-theoretic formulas (63)–(65) stated above.

Lemma 6.2 *For any $x \in \mathcal{O}(S_{qr}^2)$, we have*

$$\begin{aligned} Ax &= (K^{2\triangleright}x)A, & xA &= A(K^{-2\triangleright}x), \\ [B^*, x] &= -q^{1/2}\lambda A(K^{-1}E\triangleright x), & [B, x] &= -q^{3/2}\lambda A(K^{-1}F\triangleright x). \end{aligned}$$

In particular, the commutators $[B, x]$ and $[B^, x]$ are in $A \cdot \mathcal{O}(S_{qr}^2)$.*

It suffices to prove the lemma for elements x of the vector space basis $\{A^n B^{\#k} ; n \in \mathbb{N}_0, k \in \mathbb{Z}\}$ of $\mathcal{O}(S_{qr}^2)$. This can be done by a straightforward induction argument. We omit the details.

6.2 Description of quantum line bundles by charts

Throughout this subsection, we suppose that $j \in \frac{1}{2}\mathbb{Z}$.

From Proposition 4.6 and Theorems 3.1 and 4.1, it follows that each irreducible integrable $*$ -representation of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ can be realized as a projective module $M_j \cong \mathcal{O}(S_{qr}^2)^{2|j|+1}P_j$. It is known that the projective modules M_j can be considered as line bundles over the quantum spheres S_{qr}^2 [2, 6]. In this section, we describe the quantum line bundles M_j by two “charts”. The charts will be realized by algebras of functions on the positive and the negative part of the spectrum of the self-adjoint operator A . These function algebras lead to $*$ -representations of the $*$ -algebra $\mathcal{F}_b(S_{qr}^2) \otimes \mathcal{Y}_r$ by left and right multiplications. It will be shown that each chart is related to a tensor product of an irreducible $*$ -representation of $\mathcal{O}(S_{qr}^2)$ from Subsection 5.3 and an irreducible $*$ -representation of the $*$ -algebra $\mathcal{Y}_r$ from Subsection 5.2.

Recall that the isomorphism Ψ_j realizing $M_j \cong \mathcal{O}(S_{qr}^2)^{2|j|+1}P_j$ is given by Equation (22). In what follows, we consider $\mathcal{O}(S_{qr}^2)^{2|j|+1}P_j$ as a subspace of $\mathcal{F}_b(S_{qr}^2)^{2|j|+1}$. Our next aim is to equip $\mathcal{F}_b(S_{qr}^2)^{2|j|+1}$ with an inner product such that Ψ_j becomes an isometry. We begin by proving an auxiliary lemma.

Lemma 6.3 *Let $x \in \mathcal{O}(S_{qr}^2)$ and $k, l \in \{-|j|, -|j| + 1, \dots, -|j|\}$. Then*

$$h(v_{kj}^{[j]*} x v_{lj}^{[j]}) = c_j q^{2|j|-2k} h(x v_{lj}^{[j]} v_{kj}^{[j]*}), \quad (68)$$

where $c_j = h(v_{|j|,j}^{[j]} v_{|j|,j}^{[j]*})^{-1}$. For $y \in \mathcal{F}_b(S_{qr}^2)$ and $g \in \mathcal{F}_b(\sigma(A))$, we have

$$h(yg(A)) = h(g(A)y), \quad h(yB) = q^2 h(By), \quad h(yB^*) = q^{-2} h(B^*y). \quad (69)$$

Proof. We first show that it suffices to prove (68) for $k = |j|$. Indeed, from (20), it follows that there is a non-zero real constant γ_k^j such that $v_{kj}^{[j]} = \gamma_k^j F^{|j|-k} v_{|j|,j}^{[j]}$. Recall that $h((f \triangleright y)^* z) = h(y^* f \triangleright z)$ and $(f \triangleright y)^* = S(f)^* \triangleright y^*$ for $f \in \mathcal{U}_q(\text{su}_2)$, $y, z \in \mathcal{O}(\text{SU}_q(2))$. Assuming that (68) holds for $k = |j|$, we get

$$\begin{aligned} h(v_{kj}^{[j]*} x v_{lj}^{[j]}) &= \gamma_k^j h((F^{|j|-k} v_{|j|,j}^{[j]})^* x v_{lj}^{[j]}) = \gamma_k^j h(v_{|j|,j}^{[j]*} E^{|j|-k} (x v_{lj}^{[j]})) \\ &= c_j \gamma_k^j h((S(E^{|j|-k})^* \triangleright (x v_{lj}^{[j]})^*)^* v_{|j|,j}^{[j]*}) = c_j \gamma_k^j h(x v_{lj}^{[j]} S(E^{|j|-k}) \triangleright v_{|j|,j}^{[j]*}) \\ &= c_j \gamma_k^j h(x v_{lj}^{[j]} (S^2(E^{|j|-k})^* \triangleright v_{|j|,j}^{[j]})^*) = c_j q^{2(|j|-k)} h(x v_{lj}^{[j]} v_{kj}^{[j]*}). \end{aligned}$$

Next we note that it suffices to prove that

$$h(v_{|j|,j}^{[j]*} v_{lj}^{[j]+n}) = c_j h(v_{lj}^{[j]+n} v_{|j|,j}^{[j]*}) \quad (70)$$

for all $n \in \mathbb{N}_0$ since $x v_{lj}^{[j]} \in M_j$ and the elements $v_{lj}^{[j]+n}$ span M_j by Theorem 3.1. As the elements $v_{lj}^{[j]+n}$ form an orthonormal set in $\mathcal{O}(\text{SU}_q(2))$ with inner product defined by (16), we only have to show that the right-hand side of (70) vanishes whenever $l \neq |j|$ or $n > 0$. (Observe that Equation (70) is trivially satisfied for $l = |j|$ and $n = 0$.) If $l \neq |j|$, then it follows from $K \triangleright v_{lj}^{[j]+n} v_{|j|,j}^{[j]*} = q^{|j|-l} v_{lj}^{[j]+n} v_{|j|,j}^{[j]*}$ and the $\mathcal{U}_q(\text{su}_2)$ -invariance of h that $h(v_{lj}^{[j]+n} v_{|j|,j}^{[j]*}) = 0$. If $n > 0$, then $v_{lj}^{[j]+n} = \kappa_l^{|j|+n} F^{\triangleright} v_{l+1,j}^{[j]+n}$ with a non-zero real constant $\kappa_l^{|j|+n}$. Hence

$$\begin{aligned} h(v_{lj}^{[j]+n} v_{|j|,j}^{[j]*}) &= \kappa_l^{|j|+n} h((S(F)^* \triangleright v_{l+1,j}^{[j]+n*})^* v_{|j|,j}^{[j]*}) = \kappa_l^{|j|+n} h(v_{l+1,j}^{[j]+n*} S(F) \triangleright v_{|j|,j}^{[j]*}) \\ &= \kappa_l^{|j|+n} q^{-2} h(v_{l+1,j}^{[j]+n*} (E \triangleright v_{|j|,j}^{[j]})^*) = 0 \end{aligned}$$

since $v_{|j|,j}^{[j]}$ is a highest weight vector.

To prove (69), we can assume that $y = f(A)B^{\#k}$, where $f \in \mathcal{F}_b(\sigma(A))$ and $k \in \mathbb{Z}$, because these elements span $\mathcal{F}_b(S_{qr}^2)$. Since $h(yg(A)) = h(g(A)y) = 0$ for $k \neq 0$, the first equality of (69) is obvious. The third equality of (69) follows from the second one because the state h is hermitian. As $h(yB) = h(By) = 0$ for $k \neq -1$, it remains to treat the case $k = -1$. Using the relations $B^*B =$

$(A - \lambda_-)(\lambda_+ - A)$, $BB^* = (q^2 A - \lambda_-)(\lambda_+ - q^2 A)$ and Equation (61), we get

$$\begin{aligned} h(yB) &= h(f(A)B^*B) = h(f(A)(A - \lambda_-)(\lambda_+ - A)) \\ &= h(q^2 f(q^2 A)(q^2 A - \lambda_-)(\lambda_+ - q^2 A)) = h(q^2 f(q^2 A)BB^*) \\ &= h(q^2 Bf(A)B^*) = h(q^2 By). \end{aligned} \quad \square$$

Proposition 6.4 *Let $\mathcal{L}_2(\mathbb{S}_{qr}^2)^{2|j|+1}$ be the Hilbert space completion of $\mathcal{F}_b(\mathbb{S}_{qr}^2)^{2|j|+1}$ with respect to the inner product given by*

$$\langle (y_{-|j|}, \dots, y_{|j|}), (z_{-|j|}, \dots, z_{|j|}) \rangle = c_j q^{2|j|} \sum_{k=-|j|}^{|j|} q^{-2k} h(z_k^* y_k), \quad (71)$$

where $c_j = h(v_{|j|,j}^{[j]} v_{|j|,j}^{[j]*})^{-1}$. Then the right multiplication by P_j on $\mathcal{F}_b(\mathbb{S}_{qr}^2)^{2|j|+1}$ defines an orthogonal projection on the Hilbert space $\mathcal{L}_2(\mathbb{S}_{qr}^2)^{2|j|+1}$. The isomorphism $\Psi_j : \mathcal{O}(\mathbb{S}_{qr}^2)^{2|j|+1} P_j \xrightarrow{\cong} M_j$ from Equation (22) is an isometry.

Proof. As $P_j^2 = P_j$, we only have to show that P_j is self-adjoint with respect to the inner product (71). Since $K \triangleright v_{lj}^{[j]} v_{kj}^{[j]*} = q^{l-k} v_{lj}^{[j]} v_{kj}^{[j]*}$, there exists a polynomial $p_{lk}^j(A)$ such that $v_{lj}^{[j]} v_{kj}^{[j]*} = B^{\#k-l} p_{lk}^j(A)$. Hence, by (69),

$$h(x v_{lj}^{[j]} v_{kj}^{[j]*}) = h(q^{2(k-l)} v_{lj}^{[j]} v_{kj}^{[j]*} x) \quad (72)$$

for $x \in \mathcal{F}_b(\mathbb{S}_{qr}^2)$. This gives

$$\begin{aligned} &\langle (y_{-|j|}, \dots, y_{|j|}) P_j, (z_{-|j|}, \dots, z_{|j|}) \rangle \\ &= c_j q^{2|j|} \sum_{k,l=-|j|}^{|j|} q^{-2k} h([2|j|+1]^{-1} q^{-(l+k)} z_k^* y_l v_{lj}^{[j]} v_{kj}^{[j]*}) \\ &= c_j q^{2|j|} \sum_{k,l=-|j|}^{|j|} q^{-2l} h([2|j|+1]^{-1} q^{-(l+k)} v_{lj}^{[j]} v_{kj}^{[j]*} z_k^* y_l) \\ &= \langle (y_{-|j|}, \dots, y_{|j|}), (z_{-|j|}, \dots, z_{|j|}) P_j \rangle, \end{aligned}$$

which proves the first assertion of the proposition.

Let $(y_{-|j|}, \dots, y_{|j|}) = (y_{-|j|}, \dots, y_{|j|}) P_j \in \mathcal{O}(\mathbb{S}_{qr}^2)^{2|j|+1} P_j$ and $(z_{-|j|}, \dots, z_{|j|}) = (z_{-|j|}, \dots, z_{|j|}) P_j \in \mathcal{O}(\mathbb{S}_{qr}^2)^{2|j|+1} P_j$. From the definition of P_j , it follows that

$y_n = [2|j|+1]^{-1} \sum_{k=-|j|}^{|j|} q^{-(k+n)} y_k v_{kj}^{[j]} v_{nj}^{[j]*}$. Using Lemma 6.3 and the last identity, we compute

$$\begin{aligned}
& \langle \Psi_j(y_{-|j|}, \dots, y_{|j|}), \Psi_j(z_{-|j|}, \dots, z_{|j|}) \rangle \\
&= [2|j|+1]^{-1} h \left(\left(\sum_{n=-|j|}^{|j|} q^{-n} z_n v_{nj}^{[j]} \right)^* \sum_{k=-|j|}^{|j|} q^{-k} y_k v_{kj}^{[j]} \right) \\
&= [2|j|+1]^{-1} \sum_{n,k=-|j|}^{|j|} h(q^{-(n+k)} v_{nj}^{[j]*} z_n^* y_k v_{kj}^{[j]}) \\
&= [2|j|+1]^{-1} \sum_{n,k=-|j|}^{|j|} c_j q^{2|j|-2n} h(q^{-(n+k)} z_n^* y_k v_{kj}^{[j]} v_{nj}^{[j]*}) \\
&= c_j q^{2|j|} \sum_{n=-|j|}^{|j|} q^{-2n} h(z_n^* y_n) = \langle (y_{-|j|}, \dots, y_{|j|}), (z_{-|j|}, \dots, z_{|j|}) \rangle,
\end{aligned}$$

which shows the second assertion of the proposition. $\square$

Since Ψ_j is an isometric isomorphism, the $*$ -representation $\hat{\pi}_j$ of the cross product algebra $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ on M_j is unitarily equivalent to the $*$ -representation $\check{\pi}_j := \Psi_j^{-1} \circ \hat{\pi}_j \circ \Psi_j$ on $\mathcal{O}(S_{qr}^2)^{2|j|+1} P_j$. The restriction of $\check{\pi}_j$ to $\mathcal{O}(S_{qr}^2) \otimes \mathcal{Y}_r$ can be described by left and right multiplications by matrices with entries from $\mathcal{O}(S_{qr}^2)$. Clearly, for $z \in \mathcal{O}(S_{qr}^2)$, $\check{\pi}_j(z)(y_{-|j|}, \dots, y_{|j|}) P_j = z(y_{-|j|}, \dots, y_{|j|}) P_j$. Since $X = q^{3/2} \lambda A F K^{-1} + q^{-1} B$ commutes with $y_k \in \mathcal{O}(S_{qr}^2)$, we obtain

$$\begin{aligned}
[2|j|+1]^{1/2} \check{\pi}_j(X)(y_{-|j|}, \dots, y_{|j|}) &= \Psi_j^{-1} \left(\sum_{k=-|j|}^{|j|} q^{-k} y_k X v_{kj}^{[j]} \right) \\
&= \Psi_j^{-1} \left(\sum_{k=-|j|}^{|j|} q^{-k} y_k (q^{3/2-k} \lambda [|j|-k+1]^{1/2} [|j|+k]^{1/2} A v_{k-1,j}^{[j]} + q^{-1} B v_{kj}^{[j]}) \right).
\end{aligned}$$

This shows that $\check{\pi}_j(X)((y_{-|j|}, \dots, y_{|j|}) P_j) = ((y_{-|j|}, \dots, y_{|j|}) \mathfrak{m}_j) P_j$, where the matrix $\mathfrak{m}_j = (m_{kl}^j)_{k,l=-|j|}^{|j|} \in M_{2|j|+1}(\mathcal{O}(S_{qr}^2))$ has the entries

$$m_{kk}^j = q^{-1} B, \quad m_{k,k-1}^j = q^{1/2-k} \lambda [|j|-k+1]^{1/2} [|j|+k]^{1/2} A, \quad m_{kl}^j = 0, \quad l \neq k, k-1.$$

Similarly, we have $\check{\pi}_j(X^*)((y_{-|j|}, \dots, y_{|j|}) P_j) = ((y_{-|j|}, \dots, y_{|j|}) \mathfrak{m}_j^\dagger) P_j$ with matrix $\mathfrak{m}_j^\dagger = (m_{kl}^{\dagger j})_{k,l=-|j|}^{|j|}$ given by

$$m_{kk}^{\dagger j} = q B^*, \quad m_{k-1,k}^{\dagger j} = q^{5/2-k} \lambda [|j|-k+1]^{1/2} [|j|+k]^{1/2} A, \quad m_{kl}^{\dagger j} = 0, \quad l \neq k, k-1$$

and $\tilde{\pi}_j(Y)((y_{-|j|}, \dots, y_{|j|})P_j) = ((y_{-|j|}, \dots, y_{|j|})\mathbf{n}_j)P_j$ with $\mathbf{n}_j = (n_{kl}^j)_{k,l=-|j|}^{|j|}$ given by

$$n_{kk}^j = q^{-2k+1}A, \quad n_{kl}^j = 0, \quad l \neq k.$$

It is easy to check that

$$\mathbf{n}_j \mathbf{m}_j^\dagger = q^2 \mathbf{m}_j^\dagger \mathbf{n}_j, \quad \mathbf{n}_j \mathbf{m}_j = q^{-2} \mathbf{m}_j \mathbf{n}_j, \quad \mathbf{m}_j \mathbf{m}_j^\dagger - q^2 \mathbf{m}_j^\dagger \mathbf{m}_j = (1 - q^2)(\mathbf{n}_j^2 + r). \quad (73)$$

Note that all operators are bounded. Hence the restriction of $\tilde{\pi}_j$ to $\mathcal{O}(S_{qr}^2) \otimes \mathcal{Y}_r$ yields a bounded $*$ -representation $\tilde{\pi}_j^b$ on the Hilbert space $\mathcal{L}_2(S_{qr}^2)^{2|j|+1}P_j$. This representation can be extended to a bounded $*$ -representation, denoted again by $\tilde{\pi}_j^b$, of $\mathcal{F}_b(S_{qr}^2) \otimes \mathcal{Y}_r$ on $\mathcal{L}_2(S_{qr}^2)^{2|j|+1}P_j$ such that $\tilde{\pi}_j^b(f)$, $f \in \mathcal{F}_b(S_{qr}^2)$, acts on $\mathcal{F}_b(S_{qr}^2)^{2|j|+1}P_j$ by left multiplication.

Let χ_- and χ_+ denote the characteristic functions of the intervals $(-\infty, 0]$ and $[0, \infty)$, respectively. Set $\mathcal{F}_b(S_{qr}^2)_+ := \mathcal{F}_b(S_{qr}^2)\chi_+(A)$ and, for $r > 0$, $\mathcal{F}_b(S_{qr}^2)_- := \mathcal{F}_b(S_{qr}^2)\chi_-(A)$. Since $\chi_\pm(A)$ commutes with all elements of $\mathcal{F}_b(S_{qr}^2)$, $\mathcal{F}_b(S_{qr}^2)_\pm$ is a unital $*$ -algebra with unit element $\chi_\pm(A)$ and an ideal of $\mathcal{F}_b(S_{qr}^2)$. Our next aim is to describe the “charts” $\mathcal{F}_b(S_{qr}^2)^{2|j|+1}_\pm P_j$ of the quantum line bundle $\mathcal{F}_b(S_{qr}^2)^{2|j|+1}P_j$ by the function algebras $\mathcal{F}_b(S_{qr}^2)_-$ and $\mathcal{F}_b(S_{qr}^2)_+$ themselves.

Lemma 6.5 *Set $\xi(s) := -sab + (s^2 - 1)qbc + sq^2dc$, where the parameter s is defined as in Section 3. Then*

- (i) $(a - qsc)(d + sb) = 1 - \xi(s), \quad (d + q^{-1}sb)(a - sc) = 1 - q^{-2}\xi(s),$
 $(b - qsd)(-qc - qsa) = q^2s^2 + \xi(s), \quad (qc + sa)(-b + sd) = s^2 + \xi(s).$
- (ii) $(a - qsc)\xi(qs) = q^2\xi(s)(a - qsc), \quad (d + q^{-1}sb)\xi(q^{-1}s) = q^{-2}\xi(s)(d + q^{-1}sb),$
 $(b - qsd)\xi(qs) = \xi(s)(b - qsd), \quad (c + q^{-1}sa)\xi(q^{-1}s) = \xi(s)(c + q^{-1}sa).$
- (iii) $v_{jj}^j v_{jj}^{j*} = \gamma_j^+(\lambda_+ - A)(\lambda_+ - q^{-2}A) \dots (\lambda_+ - q^{-4j+2}A), \quad j > 0,$
 $v_{-j,j}^j v_{-j,j}^{j*} = \gamma_j^-(q^2A - \lambda_-)(q^4A - \lambda_-) \dots (q^{4j}A - \lambda_-), \quad j > 0,$
 $v_{jj}^{|j|} v_{jj}^{|j|*} = \gamma_j^+(\lambda_+ - q^2A)(\lambda_+ - q^4A) \dots (\lambda_+ - q^{4|j|}A), \quad j < 0,$
 $v_{|j|,j}^{|j|} v_{|j|,j}^{|j|*} = \gamma_j^-(A - \lambda_-)(q^{-2}A - \lambda_-) \dots (q^{-4|j|+2}A - \lambda_-), \quad j < 0,$
with non-zero constants $\gamma_j^\pm \in \mathbb{R}$.
- (iv) *The function $v_{jj}^{|j|} v_{jj}^{|j|*} \chi_-(A)$ is invertible in $\mathcal{F}_b(S_{qr}^2)_-$ for $r > 0$.*
The function $v_{-j,j}^{|j|} v_{-j,j}^{|j|} \chi_+(A)$ is invertible in $\mathcal{F}_b(S_{qr}^2)_+$.*

$$(v) \quad v_{kj}^{|j|} v_{jj}^{|j|*} (v_{jj}^{|j|} v_{jj}^{|j|*} \chi_-(A))^{-1} v_{jj}^{|j|} v_{lj}^{|j|*} = v_{kj}^{|j|} v_{lj}^{|j|*} \chi_-(A) \text{ for } r > 0,$$

$$v_{kj}^{|j|} v_{-j,j}^{|j|*} (v_{-j,j}^{|j|} v_{-j,j}^{|j|*} \chi_+(A))^{-1} v_{-j,j}^{|j|} v_{lj}^{|j|*} = v_{kj}^{|j|} v_{lj}^{|j|*} \chi_+(A).$$

Proof. (i) and (ii) follow by straightforward computations.

(iii) Let $j > 0$. Then $v_{jj}^j = N_{jj}^j u_j$, where u_j is defined by Equation (17). From (i) and (ii), it follows that

$$(d+q^{-1}sb)\dots(d+q^{-2j}sb)(a-q^{-2j+1}sc)\dots(a-sc) = (1-q^{-2}\xi(s))\dots(1-q^{-4j}\xi(s)).$$

Inserting $\xi(s) = sr^{-1/2}q^2A = \lambda_+^{-1}q^2A$ gives the result. The other cases are treated analogously.

(iv) As $A < 0$ on the interval $(-\infty, 0]$ and $\lambda_+ > 0$, each factor $(\lambda_+ - q^{2k}A)$, $k \in \mathbb{Z}$, is invertible in $\mathcal{F}_b(S_{qr}^2)_-$. Likewise, for $r > 0$, each factor $(q^{2k}A - \lambda_-)$, $k \in \mathbb{Z}$, is invertible in $\mathcal{F}_b(S_{qr}^2)_+$ since $\lambda_- < 0$.

(v) Let $j > 0$. In the proof of Proposition 6.4, we argued that $v_{kj}^j v_{jj}^{j*} = B^{j-k} p_{kj}^j(A)$. Note that $B\xi(s) = q^2\xi(s)B$. This together with (iii) gives

$$v_{kj}^j v_{jj}^{j*} (v_{jj}^j v_{jj}^{j*} \chi_-(A))^{-1} v_{jj}^j v_{lj}^{j*} \\ = (N_{jj}^{j2} (1 - q^{2(j-k-1)}\xi(s)) \dots (1 - q^{-2(j+k)}\xi(s)) \chi_-(A))^{-1} v_{kj}^j v_{jj}^{j*} v_{jj}^j v_{lj}^{j*}.$$

On the other hand, $v_{jj}^{j*} v_{jj}^j = N_{jj}^{j2} (1 - \xi(q^{-2j}s)) \dots (1 - q^{4j-2}\xi(q^{-2j}s))$ by (i) and (ii). The vector v_{kj}^j is a linear combination of products consisting of factors $(d+q^{-n}sb)$ and $(c+q^{-m}sa)$, $1 \leq n, m \leq 2j$, where the terms $(d+q^{-n}sb)$ and $(c+q^{-m}sa)$ appear $j+k$ -times and $j-k$ -times, respectively. Applying (i) and (ii), we see that $v_{kj}^j v_{jj}^{j*} v_{jj}^j = N_{jj}^{j2} (1 - q^{-2(j+k)}\xi(s)) \dots (1 - q^{2(j-k-1)}\xi(s)) v_{kj}^j$. Inserting this identity into above equation gives the result. The other cases are handled similarly. $\square$

Lemma 6.6 *With the inner product on $\mathcal{F}_b(S_{qr}^2)_\pm$ defined by $\langle f, g \rangle := h(g^*f)$, there is an isometric isomorphism $\check{\Psi}_{j,\pm}$ from $\mathcal{F}_b(S_{qr}^2)_\pm$ onto $\mathcal{F}_b(S_{qr}^2)_\pm^{2|j|+1} P_j$ given by*

$$\begin{aligned} \check{\Psi}_{j,-}(f) &= q^{2j}[2j+1]^{1/2} (0, \dots, 0, f(v_{jj}^j v_{jj}^{j*} \chi_-(A))^{-1/2}) P_j, \quad j > 0, \\ \check{\Psi}_{j,-}(f) &= q^{2j}[2|j|+1]^{1/2} (f(v_{jj}^{|j|} v_{jj}^{|j|*} \chi_-(A))^{-1/2}, 0, \dots, 0) P_j, \quad j < 0, \\ \check{\Psi}_{j,+}(g) &= q^{-2j}[2j+1]^{1/2} (g(v_{-j,j}^j v_{-j,j}^{j*} \chi_+(A))^{-1/2}, 0, \dots, 0) P_j, \quad j > 0, \\ \check{\Psi}_{j,+}(g) &= q^{-2j}[2|j|+1]^{1/2} (0, \dots, 0, g(v_{|j|,j}^{|j|} v_{|j|,j}^{|j|*} \chi_+(A))^{-1/2}) P_j, \quad j < 0, \end{aligned}$$

where $f \in \mathcal{F}_b(S_{qr}^2)_-$ and $g \in \mathcal{F}_b(S_{qr}^2)_+$.

Proof. We carry out the proof for $\check{\Psi}_{j,-}$ and $j > 0$. The other cases are treated in the same manner. By Lemma 6.5, $v_{jj}^j v_{jj}^{j*} \chi_-(A)$ is invertible. Hence $\check{\Psi}_{j,-}$ is well defined. Fix $(y_{-j}, \dots, y_j) = (y_{-j}, \dots, y_j) P_j \in \mathcal{F}_b(S_{qr}^2)_{-}^{2j+1} P_j$. Let $z = \sum_{k=-j}^j q^{j-k} y_k v_{kj}^j v_{jj}^{j*} (v_{jj}^j v_{jj}^{j*} \chi_-(A))^{-1}$ and $(z_{-j}, \dots, z_j) = (0, \dots, 0, z) P_j$. By Lemma 6.5,

$$\begin{aligned} z_l &= [2j+1]^{-1} \sum_{k=-j}^j q^{-(k+l)} y_k v_{kj}^j v_{jj}^{j*} (v_{jj}^j v_{jj}^{j*} \chi_-(A))^{-1} v_{jj}^j v_{lj}^{j*} \\ &= [2j+1]^{-1} \sum_{k=-j}^j q^{-(k+l)} y_k v_{kj}^j v_{lj}^{j*} \chi_-(A) = y_l. \end{aligned}$$

Thus $\check{\Psi}_{j,-}(q^{-2j} [2j+1]^{-1/2} z (v_{jj}^j v_{jj}^{j*} \chi_-(A))^{1/2}) = (y_{-j}, \dots, y_j)$, so $\check{\Psi}_{j,-}$ is surjective.

Next we verify that $\check{\Psi}_{j,-}$ is isometric. Let $f \in \mathcal{F}_b(S_{qr}^2)_{-}$. Since P_j is a projection, we have $[2j+1]^{-2} \sum_{k=-j}^j q^{-2(j+k)} v_{jj}^j v_{kj}^{j*} v_{kj}^j v_{jj}^{j*} = [2j+1]^{-1} q^{-2j} v_{jj}^j v_{jj}^{j*}$. Using Equations (69) and (72), we conclude that

$$h(x(v_{jj}^j v_{jj}^{j*} \chi_-(A))^{-1/2} v_{jj}^j v_{kj}^{j*}) = q^{2(k-j)} h((v_{jj}^j v_{jj}^{j*} \chi_-(A))^{-1/2} v_{jj}^j v_{kj}^{j*} x).$$

From these relations, it follows that, for $f \in \mathcal{F}_b(S_{qr}^2)_{-}$,

$$\begin{aligned} \|\check{\Psi}_{j,-}(f)\|^2 &= [2j+1]^{-1} q^{4j} \sum_{k=-j}^j q^{-2k} \\ &\quad \times h(q^{-2(j+k)} v_{jj}^j v_{kj}^{j*} (v_{jj}^j v_{jj}^{j*} \chi_-(A))^{-1/2} f^* f (v_{jj}^j v_{jj}^{j*} \chi_-(A))^{-1/2} v_{jj}^j v_{kj}^{j*}) \\ &= h(f^* f) = \|f\|^2. \end{aligned} \quad \square$$

As multiplication by elements of $\mathcal{F}_b(S_{qr}^2)$ leaves the decomposition $\mathcal{F}_b(S_{qr}^2) = \mathcal{F}_b(S_{qr}^2)_{-} \oplus \mathcal{F}_b(S_{qr}^2)_{+}$ invariant, we have $\mathcal{F}_b(S_{qr}^2)^{2|j|+1} P_j = \mathcal{F}_b(S_{qr}^2)_{-}^{2|j|+1} P_j \oplus \mathcal{F}_b(S_{qr}^2)_{+}^{2|j|+1} P_j$ and the $*$ -representation $\check{\pi}_j^b$ of $\mathcal{F}_b(S_{qr}^2) \otimes \mathcal{Y}_r$ decomposes into a direct sum $\check{\pi}_j^b = \check{\pi}_{j,-}^b \oplus \check{\pi}_{j,+}^b$ of $*$ -representations $\check{\pi}_{j,\pm}^b$ on $\mathcal{F}_b(S_{qr}^2)_{\pm}^{2|j|+1} P_j$. Using the isometric isomorphism $\check{\Psi}_{j,\pm}$, the $*$ -representation $\check{\pi}_{j,\pm}^b$ is unitarily equivalent to a $*$ -representation $\rho_{j,\pm} := (\check{\Psi}_{j,\pm})^{-1} \circ \check{\pi}_{j,\pm}^b \circ \check{\Psi}_{j,\pm}$ of $\mathcal{F}_b(S_{qr}^2) \otimes \mathcal{Y}_r$ on $\mathcal{F}_b(S_{qr}^2)_{\pm}$.

Theorem 6.7 *The $*$ -representation $\rho_{j,\pm}$ of $\mathcal{F}_b(S_{qr}^2) \otimes \mathcal{Y}_r$ on $\mathcal{F}_b(S_{qr}^2)_{\pm}$ is given by*

$$\begin{aligned} \rho_{j,-}(X)(f) &= q^{-1} f B(\lambda_+ - q^{-4j} A)^{1/2} (\lambda_+ - A)^{-1/2}, \quad \rho_{j,-}(Y)(f) = q^{-2j+1} f A, \\ \rho_{j,-}(X^*)(f) &= q f (\lambda_+ - q^{-4j} A)^{1/2} (\lambda_+ - A)^{-1/2} B^*, \end{aligned}$$

$$\begin{aligned}\rho_{j,+}(X)(f) &= q^{-1}fB(q^{4j}A - \lambda_-)^{1/2}(A - \lambda_-)^{-1/2}, \quad \rho_{j,+}(Y)(f) = q^{2j+1}fA, \\ \rho_{j,+}(X^*)(f) &= qf(q^{4j}A - \lambda_-)^{1/2}(A - \lambda_-)^{-1/2}B^*,\end{aligned}$$

and $\rho_{j,\pm}(x)(f) = xf$ for $x \in \mathcal{F}_b(S_{qr}^2)$, $f \in \mathcal{F}_b(S_{qr}^2)_{\pm}$.

In particular, all representation operators $\rho_{j,\pm}(y)$, $y \in \mathcal{F}_b(S_{qr}^2) \otimes \mathcal{Y}_r$, are bounded and $\rho_{j,\pm}$ extends to a $*$ -representation, denoted also by $\rho_{j,\pm}$, on the Hilbert space completion $\mathcal{L}_2(S_{qr}^2)_{\pm}$ of $\mathcal{F}_b(S_{qr}^2)_{\pm}$. The restriction of this $*$ -representation $\rho_{j,\pm}$ to $\mathcal{O}(S_{qr}^2) \otimes \mathcal{Y}_r$ is unitarily equivalent to a tensor product representation $\sigma^{\pm} \otimes \sigma_j^{\pm}$ on $\mathcal{H}^{\pm} \otimes \mathcal{K}^{\pm}$, where $\sigma^{\pm}$ denotes the irreducible $*$ -representation of $\mathcal{O}(S_{qr}^2)$ on $\mathcal{H}^{\pm}$ from Subsection 5.3 and $\sigma_j^{\pm}$ denotes the irreducible $*$ -representation of $\mathcal{Y}_r$ on $\mathcal{K}^{\pm}$ from Subsection 5.2 with $Y_0 = q^{\pm 2j+1}\lambda_{\pm}$. The restriction of the $*$ -representation $\check{\pi}_j \cong \hat{\pi}_j \cong \pi_j$ of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ to $\mathcal{O}(S_{qr}^2) \otimes \mathcal{Y}_r$ is unitarily equivalent to the direct sum representation $\rho_{j,-} \oplus \rho_{j,+}$.

Proof. Clearly, for $x \in \mathcal{F}_b(S_{qr}^2)$, $\rho_{j,-}(x)$ acts by left multiplication, that is, $\rho_{j,-}(x)(g) = xg$, $g \in \mathcal{F}_b(S_{qr}^2)_{-}$. Our next aim is to compute the action of the generators of $\mathcal{Y}_r$. Let $g \in \mathcal{F}_b(S_{qr}^2)_{-}$. By the definition the matrix $\mathfrak{n}_j$,

$$\begin{aligned}\rho_{j,-}(Y)(g) &= (\check{\Psi}_{j,-})^{-1}(q^{2j}[2j+1]^{1/2}(0, \dots, 0, f(v_{jj}^j v_{jj}^{j*} \chi_{-}(A))^{-1/2}) \mathfrak{n}_j P_j) \\ &= (\check{\Psi}_{j,-})^{-1}(q^{2j}[2j+1]^{1/2}(0, \dots, 0, q^{-2j+1}fA(v_{jj}^j v_{jj}^{j*} \chi_{-}(A))^{-1/2}) P_j) = q^{-2j+1}fA.\end{aligned}$$

From Lemma 6.5 (iii), it follows that

$$(v_{jj}^j v_{jj}^{j*} \chi_{-}(A))^{-1/2} B^* (v_{jj}^j v_{jj}^{j*} \chi_{-}(A))^{1/2} = \chi_{-}(A) (\lambda_{+} - q^{-4j} A)^{1/2} (\lambda_{+} - A)^{-1/2} B^*.$$

Using this identity and the explicit form of the matrix $\mathfrak{m}_j^{\dagger}$, we see that

$$\begin{aligned}\rho_{j,-}(X^*)(g) &= (\check{\Psi}_{j,-})^{-1}(q^{2j}[2j+1]^{1/2}(0, \dots, 0, f(v_{jj}^j v_{jj}^{j*} \chi_{-}(A))^{-1/2}) \mathfrak{m}_j^{\dagger} P_j) \\ &= qg(\lambda_{+} - q^{-4j} A)^{1/2} (\lambda_{+} - A)^{-1/2} B^*.\end{aligned}$$

The operator $\rho_{j,-}(X)$ is determined by the relation $\rho_{j,-}(X) = \rho_{j,-}(X^*)^*$. Since $h(g^* f \varphi(A) B^*) = h(q^{-2}(gB\bar{\varphi}(A))^* f)$ for all $f, g, \varphi(A) \in \mathcal{F}_b(S_{qr}^2)_{-}$, we obtain

$$\rho_{j,-}(X)(g) = q^{-1}gB(\lambda_{+} - q^{-4j} A)^{1/2} (\lambda_{+} - A)^{-1/2}.$$

This proves the formulas of the theorem for $\rho_{j,-}$, $j > 0$. The other cases are treated in the same way. From the preceding formulas it is clear that all representation operators $\rho_{j,\pm}(y)$, $y \in \mathcal{F}_b(S_{qr}^2) \otimes \mathcal{Y}_r$, are bounded.

For $n \in \mathbb{N}_0$, let $\chi_{n,\pm}$ denote the characteristic function of the point $q^{2n}\lambda_{\pm}$. Define

$$\vartheta_{nl}^{\pm} := c_{nl}^{\pm} \chi_{n,\pm}(A) B^{\#l}, \quad n \in \mathbb{N}_0, \quad l \in \mathbb{Z}, \quad l \geq -n, \quad (74)$$

where, with $\gamma_{\pm}$ defined in Subsection 6.1,

$$\begin{aligned} c_{n,0}^{\pm} &= \gamma_{\pm}^{-1/2} q^{-n}, \\ c_{nl}^{\pm} &= \gamma_{\pm}^{-1/2} q^{-n-l} \left(\prod_{m=0}^{|l|-1} (q^{2(n-m)} \lambda_{\pm} - \lambda_{\mp})(\lambda_{+} - q^{2(n-m)} \lambda_{\pm}) \right)^{-1/2}, \quad l < 0, \\ c_{nl}^{\pm} &= \gamma_{\pm}^{-1/2} q^{-n-l} \left(\prod_{m=1}^l (q^{2(n+m)} \lambda_{\pm} - \lambda_{\mp})(\lambda_{+} - q^{2(n+m)} \lambda_{\pm}) \right)^{-1/2}, \quad l > 0. \end{aligned}$$

For $r = 0$, only $\vartheta_{nl}^{\pm}$ is considered. The set $\{\vartheta_{nl}^{\pm}; n \in \mathbb{N}_0, l \in \mathbb{Z}, l \geq -n, \}$ is an orthonormal basis of $\mathcal{L}_2(S_{qr}^2)_{\pm}$. Note that $c_{\pm}(n) = \lambda_n(q^{2n}\lambda_{\pm}^2 + r)^{1/2} = ((q^{2n}\lambda_{\pm} - \lambda_{\mp})(\lambda_{+} - q^{2n}\lambda_{\pm}))^{1/2}$. Using the commutation rules in $\mathcal{F}_b(S_{qr}^2)$ and applying $f(A)\chi_{n,\pm}(A) = f(q^{2n}\lambda_{\pm})\chi_{n,\pm}(A)$ for all $f \in \mathcal{F}_b(\sigma(A))$, we get

$$\begin{aligned} \rho_{j,\pm}(A)\vartheta_{nl}^{\pm} &= \lambda_{\pm} q^{2n} \vartheta_{nl}^{\pm}, \\ \rho_{j,\pm}(B)\vartheta_{nl}^{\pm} &= c_{\pm}(n) \vartheta_{n-1,l+1}^{\pm}, \quad \rho_{j,\pm}(B^*)\vartheta_{nl}^{\pm} = c_{\pm}(n+1) \vartheta_{n+1,l-1}^{\pm}, \\ \rho_{j,\pm}(Y)\vartheta_{nl}^{\pm} &= \lambda_{\pm} q^{\pm 2j+1} q^{2(n+l)} \vartheta_{nl}^{\pm}, \\ \rho_{j,\pm}(X)\vartheta_{nl}^{\pm} &= \lambda_{n+l+1} (q^{2(n+l)} (\lambda_{\pm} q^{\pm 2j+1})^2 + r)^{1/2} \vartheta_{n,l+1}^{\pm}, \\ \rho_{j,\pm}(X^*)\vartheta_{nl}^{\pm} &= \lambda_{n+l} (q^{2(n+l-1)} (\lambda_{\pm} q^{\pm 2j+1})^2 + r)^{1/2} \vartheta_{n,l-1}^{\pm}. \end{aligned}$$

Renaming $\zeta_{nk}^{\pm} := \vartheta_{n,k-n}^{\pm}$, $k, n \in \mathbb{N}_0$, we obtain

$$\begin{aligned} \rho_{j,\pm}(A)\zeta_{nk}^{\pm} &= \lambda_{\pm} q^{2n} \zeta_{nk}^{\pm}, \quad \rho_{j,\pm}(B)\zeta_{nk}^{\pm} = c_{\pm}(n) \zeta_{n-1,k}^{\pm}, \\ \rho_{j,\pm}(B^*)\zeta_{nk}^{\pm} &= c_{\pm}(n+1) \zeta_{n+1,k}^{\pm}, \\ \rho_{j,\pm}(Y)\zeta_{nk}^{\pm} &= q^{2k} q^{\pm 2j+1} \lambda_{\pm} \zeta_{nk}^{\pm}, \quad \rho_{j,\pm}(X)\zeta_{nk}^{\pm} = \lambda_{k+1} (q^{2k} (q^{\pm 2j+1} \lambda_{\pm})^2 + r)^{1/2} \zeta_{n,k+1}^{\pm}, \\ \rho_{j,\pm}(X^*)\zeta_{nk}^{\pm} &= \lambda_k (q^{2(k-1)} (q^{\pm 2j+1} \lambda_{\pm})^2 + r)^{1/2} \zeta_{n,k-1}^{\pm}. \end{aligned}$$

Let $\mathcal{H}^{\pm}$ and $\mathcal{K}^{\pm}$ be the closed linear spans of orthonormal systems $\{\eta_n^{\pm}; n \in \mathbb{N}_0\}$ and $\{\xi_k^{\pm}; k \in \mathbb{N}_0\}$, respectively. Setting $\zeta_{nk}^{\pm} := \eta_n^{\pm} \otimes \xi_k^{\pm}$, we see that the restriction of $\rho_{j,\pm}$ to $\mathcal{O}(S_{qr}^2) \otimes \mathcal{Y}_r$ is unitarily equivalent to the tensor product representation $\sigma^{\pm} \otimes \sigma_j^{\pm}$. The last assertion of the theorem follows immediately from the preceding since $\mathcal{O}(S_{qr}^2) \otimes \mathcal{Y}_r \subset \mathcal{F}_b(S_{qr}^2) \otimes \mathcal{Y}_r$. $\square$

7 Description of the irreducible integrable representations by the second approach

As we have seen in Subsection 6.2, the irreducible integrable representation $\tilde{\pi}_j$ of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ leads to bounded $*$ -representations $\rho_{j,\pm}$ of $\mathcal{O}(S_{qr}^2) \otimes \mathcal{Y}_r$ on the charts $\mathcal{F}_b(S_{qr}^2)_\pm$. In this section, we recover the irreducible integrable representations $\tilde{\pi}_j$ of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ by taking the orthogonal sum of both charts and passing to another domain. Because this construction is based on $*$ -representations of $\mathcal{O}(S_{qr}^2) \otimes \mathcal{Y}_r$, we say that we have described the irreducible integrable representation $\tilde{\pi}_j$ by the second approach.

We begin by showing that the bounded $*$ -representation $\rho_{j,\pm}$ of $\mathcal{O}(S_{qr}^2) \otimes \mathcal{Y}_r$ on the Hilbert space completion $\mathcal{L}_2(S_{qr}^2)_\pm$ of $\mathcal{F}_b(S_{qr}^2)_\pm$ leads to a $*$ -representation of the $*$ -algebra $\hat{\mathcal{O}}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$. By a slight abuse of notation, we use the same symbol $\rho_{j,\pm}$ to denote the representation of the cross product algebra. It is a $*$ -representation by unbounded operators acting on the invariant dense domain

$$\mathcal{D}_{j,\pm} := \cap_{n,m=0}^{\infty} \mathcal{D}(\rho_{j,\pm}(A)^{-n} \rho_{j,\pm}(Y)^{-m}) \subset \mathcal{L}_2(S_{qr}^2)_\pm$$

Note that $\pi(A)$ and $\pi(Y)$ are commuting bounded self-adjoint operators but their inverses are unbounded. For $\varphi \in \mathcal{D}_{j,\pm}$, define

$$\rho_{j,\pm}(K)\varphi := q^{1/2}|\rho_{j,\pm}(Y)|^{-1/2}|\rho_{j,\pm}(A)|^{1/2}\varphi, \quad \rho_{j,\pm}(A^{-1})\varphi := \rho_{j,\pm}(A)^{-1}\varphi \quad (75)$$

and $\rho_{j,\pm}(K^{-1}) := \rho_{j,\pm}(K)^{-1}$. When φ belongs to $\mathcal{F}_b(S_{qr}^2)_\pm \cap \mathcal{D}_{j,\pm}$, we can write $\rho_{j,\pm}(K)\varphi = q^j|A|^{1/2}\varphi|A|^{-1/2}$ and $\rho_{j,\pm}(A^{-1})\varphi := A^{-1}\varphi$. From the commutation rules in the algebra $\hat{\mathcal{O}}(S_{qr}^2) \otimes \mathcal{Y}_r$, it follows easily that (75) defines indeed a $*$ -representation of the larger $*$ -algebra $\hat{\mathcal{O}}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ on $\mathcal{D}_{j,\pm}$.

Using the isometric isomorphism $\check{\Psi}_{j,\pm}$, we obtain an irreducible $*$ -representation $\check{\rho}_{j,\pm} := \check{\Psi}_{j,\pm} \circ \rho_{j,\pm} \circ (\check{\Psi}_{j,\pm})^{-1}$ of $\hat{\mathcal{O}}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ on

$$\mathcal{D}(\check{\rho}_{j,\pm}) := \cap_{n,m=0}^{\infty} \mathcal{D}(\check{\rho}_{j,\pm}(A)^{-n} \check{\rho}_{j,\pm}(Y)^{-m}) \subset \mathcal{L}_2(S_{qr}^2)_\pm^{2|j|+1} P_j,$$

The restriction of the direct sum $\check{\rho}_j := \check{\rho}_{j,-} \oplus \check{\rho}_{j,+}$ to $\mathcal{O}(S_{qr}^2) \otimes \mathcal{Y}_r$ yields a bounded representation which can be extended to a representation on the Hilbert space $\mathcal{L}_2(S_{qr}^2)^{2|j|+1} P_j = \mathcal{L}_2(S_{qr}^2)_-^{2|j|+1} P_j \oplus \mathcal{L}_2(S_{qr}^2)_+^{2|j|+1} P_j$. By the definitions of $\rho_{j,\pm}$ and $\check{\rho}_{j,\pm}$, it is obvious that this representation coincides with $\tilde{\pi}_j^b$ and so it coincides with the restriction of $\tilde{\pi}_j$ to $\mathcal{O}(S_{qr}^2) \otimes \mathcal{Y}_r$ on its common domain. However, the restriction of $\check{\rho}_j$ to $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ is not unitarily equivalent to $\tilde{\pi}_j$ since the

latter is irreducible while the former is not. Theorem 7.2 below shows that we can reconstruct the irreducible integrable representations $\check{\pi}_j \cong \hat{\pi}_j \cong \pi_j$ from $\check{\rho}_j$. The proof of Theorem 7.2 is based on the following lemma.

Lemma 7.1 *Let $\eta \in \mathcal{O}(S_{qr}^2)^{2|j|+1}P_j$. Then*

$$\check{\pi}_j^b(A)\check{\pi}_j(K^{-1}E)\eta = q^{-3/2}\lambda^{-1}\check{\pi}_j^b(X^* - qB^*)\eta, \quad (76)$$

$$\check{\pi}_j^b(A)\check{\pi}_j(FK^{-1})\eta = q^{-3/2}\lambda^{-1}\check{\pi}_j^b(X - q^{-1}B)\eta, \quad (77)$$

$$|\check{\pi}_j^b(Y)|^{1/2}\check{\pi}_j(K)\eta = q^{1/2}|\check{\pi}_j^b(A)|^{1/2}\eta, \quad |\check{\pi}_j^b(A)|^{1/2}\check{\pi}_j(K^{-1})\eta = q^{-1/2}|\check{\pi}_j^b(Y)|^{1/2}\eta. \quad (78)$$

Proof. Let $\eta = (y_{-|j|}, \dots, y_{|j|})P_j \in \mathcal{O}(S_{qr}^2)^{2|j|+1}P_j$. From the uniqueness of the square root of a positive operator, it follows that

$$\begin{aligned} |\check{\pi}_j^b(A)|^{1/2}(y_{-|j|}, \dots, y_{|j|})P_j &= (|A|^{1/2}y_{-|j|}, \dots, |A|^{1/2}y_{|j|})P_j, \\ |\check{\pi}_j^b(Y)|^{1/2}(y_{-|j|}, \dots, y_{|j|})P_j &= q^{1/2}(q^{|j|}y_{-|j|}|A|^{1/2}, \dots, q^{-|j|}y_{|j|}|A|^{1/2})P_j. \end{aligned}$$

Further, the commutation rules in $\mathcal{F}_b(S_{qr}^2)$ imply for all $y \in \mathcal{O}(S_{qr}^2)$

$$|A|^{1/2}y = (K \triangleright y)|A|^{1/2}, \quad y|A|^{1/2} = |A|^{1/2}(K^{-1} \triangleright y).$$

On the other hand, since $\check{\pi}_j := \Psi_j^{-1} \circ \hat{\pi}_j \circ \Psi_j$, we have

$$\begin{aligned} \check{\pi}_j(K^{\pm 1})(y_{-|j|}, \dots, y_{|j|})P_j &= [2|j|+1]^{-1/2}\Psi_j^{-1}(K^{\pm 1} \triangleright \sum_{k=-|j|}^{|j|} q^{-k}y_kv_{kj}^{|j|}) \\ &= (q^{\mp|j|}K^{\pm 1} \triangleright y_{-|j|}, \dots, q^{\pm|j|}K^{\pm 1} \triangleright y_{|j|})P_j. \end{aligned}$$

Combining the preceding equations proves (78).

Computing the action of $\check{\pi}_j(FK^{-1}) = \Psi_j^{-1} \circ \hat{\pi}_j \circ \Psi_j(FK^{-1})$ on the element $\eta = (y_{-|j|}, \dots, y_{|j|})P_j$ gives

$$\begin{aligned} \check{\pi}_j(FK^{-1})(y_{-|j|}, \dots, y_{|j|})P_j &= [2|j|+1]^{-1/2}\Psi_j^{-1} \sum_{k=-|j|}^{|j|} q^{-k}((FK^{-1} \triangleright y_k)v_{kj}^{|j|} \\ &\quad + q^{-k}[|j|-k+1]^{1/2}[|j|+k]^{1/2}(K^{-2} \triangleright y_k)v_{k-1,j}^{|j|}) \end{aligned}$$

so that, by using Lemma 6.2,

$$\begin{aligned} \check{\pi}_j^b(A)\check{\pi}_j(FK^{-1})(y_{-|j|}, \dots, y_{|j|})P_j &= (A(FK^{-1} \triangleright y_{-|j|}), \dots, A(FK^{-1} \triangleright y_{|j|}))P_j \\ &\quad + q^{-2}(q^{|j|}([2|j|][1])^{1/2}A(K^{-2} \triangleright y_{-|j|+1}), \dots, q^{-|j|+1}([1][2|j|])^{1/2}A(K^{-2} \triangleright y_{|j|}), 0)P_j \\ &= -q^{-5/2}\lambda^{-1}([B, y_{-|j|}], \dots, [B, y_{|j|}])P_j \\ &\quad + q^{-2}(q^{|j|}([2|j|][1])^{1/2}y_{-|j|+1}A, \dots, q^{-|j|+1}([1][2|j|])^{1/2}y_{|j|}A, 0)P_j. \end{aligned}$$

Comparing the last identity with the action of $\tilde{\pi}_j^b(X)$ and $\tilde{\pi}_j^b(B)$ on $\mathcal{O}(S_{qr}^2)^{2|j|+1}P_j$ from Subsection 6.2 (see the discussion preceding Proposition 6.4) shows (77). Equation (76) is proved similarly. $\square$

Let us recall the notion of the adjoint of a $*$ -representation π of a $*$ -algebra $\mathcal{X}$ (see e.g. [14, Section 8.1]). It is a representation π^* of $\mathcal{X}$ acting on the domain $\mathcal{D}(\pi^*) := \cap_{x \in \mathcal{X}} \mathcal{D}(\pi(x)^*)$ by $\pi^*(x)\varphi = \pi(x^*)^*\varphi$, where $\varphi \in \mathcal{D}(\pi^*)$. In general, π^* is not a $*$ -representation.

Theorem 7.2 *The irreducible integrable $*$ -representation $\tilde{\pi}_j$ of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ is the restriction of the adjoint $\check{\rho}_j^*$ of the $*$ -representation $\check{\rho}_j = \check{\rho}_{j,-} \oplus \check{\rho}_{j,+}$ of $\hat{\mathcal{O}}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ to the domain $\mathcal{O}(S_{qr}^2)^{2|j|+1}P_j$. There exists a Hilbert space basis $\{\eta_{nm}^\pm; n, m \in \mathbb{N}_0\}$ of $\mathcal{L}_2(S_{qr}^2)_{\pm}^{2|j|+1}P_j$ such that $\check{\rho}_{j,\pm}$ is determined by the formulas $(I)_{\pm, q \mp j}$ from Subsection 5.3.*

Proof. First note that, by definition, $\check{\rho}_j(Z)\varphi = \tilde{\pi}_j^b(Z)\varphi$ for all $Z \in \mathcal{O}(S_{qr}^2) \otimes \mathcal{Y}_r$ and $\varphi \in \mathcal{D}(\check{\rho}_{j,\pm})$. In particular, $\langle \check{\rho}_j(A^{-1})\varphi, \tilde{\pi}_j^b(A)\eta \rangle = \langle \varphi, \eta \rangle$ for all η from the Hilbert space $\mathcal{L}_2(S_{qr}^2)^{2|j|+1}P_j$. Now let $\eta \in \mathcal{O}(S_{qr}^2)^{2|j|+1}P_j$ and $\varphi \in \mathcal{D}(\check{\rho}_{j,\pm})$. Then, by Lemma 7.1,

$$\begin{aligned} \langle \check{\rho}_j(K^{-1}E)\varphi, \eta \rangle &= q^{-3/2}\lambda^{-1} \langle \check{\rho}_j(X^* - q^{-1}B^*)\check{\rho}_j(A^{-1})\varphi, \eta \rangle \\ &= q^{-3/2}\lambda^{-1} \langle \check{\rho}_j(A^{-1})\varphi, \tilde{\pi}_j^b(X - q^{-1}B)\eta \rangle \\ &= \langle \check{\rho}_j(A^{-1})\varphi, \tilde{\pi}_j^b(A)\tilde{\pi}_j(FK^{-1})\eta \rangle = \langle \varphi, \tilde{\pi}_j(FK^{-1})\eta \rangle. \end{aligned}$$

Similarly one shows $\langle \check{\rho}_j(FK^{-1})\varphi, \eta \rangle = \langle \varphi, \tilde{\pi}_j(K^{-1}E)\eta \rangle$. As above, we have $\langle |\check{\rho}_j(Y)|^{-1/2}\varphi, |\tilde{\pi}_j^b(Y)|^{1/2}\eta \rangle = \langle \varphi, \eta \rangle$ and thus, again by Lemma 7.1,

$$\begin{aligned} \langle \check{\rho}_j(K)\varphi, \eta \rangle &= q^{1/2} \langle |\check{\rho}_j(A)|^{1/2}|\check{\rho}_j(Y)|^{-1/2}\varphi, \eta \rangle = q^{1/2} \langle |\check{\rho}_j(Y)|^{-1/2}\varphi, |\tilde{\pi}_j^b(A)|^{1/2}\eta \rangle \\ &= \langle |\check{\rho}_j(Y)|^{-1/2}\varphi, |\tilde{\pi}_j^b(Y)|^{1/2}\tilde{\pi}_j(K)\eta \rangle = \langle \varphi, \tilde{\pi}_j(K)\eta \rangle. \end{aligned}$$

Likewise, $\langle \check{\rho}_j(K^{-1})\varphi, \eta \rangle = \langle \varphi, \tilde{\pi}_j(K^{-1})\eta \rangle$. Clearly, $\langle \check{\rho}_j(x^*)\varphi, \eta \rangle = \langle \varphi, \tilde{\pi}_j(x)\eta \rangle$ for all $x \in \mathcal{O}(S_{qr}^2)$. As the elements $K^{\pm 1}$, $K^{-1}E$, FK^{-1} and $x \in \mathcal{O}(S_{qr}^2)$ generate the algebra $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$, it follows that $\mathcal{O}(S_{qr}^2)^{2|j|+1}P_j$ is contained in $\mathcal{D}(\check{\rho}_j^*)$ and $\check{\rho}_j^*(Z)\eta = \tilde{\pi}_j(Z)\eta$ for all $Z \in \mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ and $\eta \in \mathcal{O}(S_{qr}^2)^{2|j|+1}P_j$.

Computing the action of $\rho_{j,\pm}(K)$, $\rho_{j,\pm}(E) = q^{-3/2}\lambda^{-1}\rho_{j,\pm}(A^{-1}K(X^* - qB^*))$, $\rho_{j,\pm}(F) = \rho_{j,\pm}(E)^*$ on the basis vectors $\zeta_{nm}^\pm$ from the proof of Theorem 6.7 shows that $\rho_{j,\pm}$ is determined by the formulas $(I)_{\pm, q \mp j}$ from Subsection 5.3. Setting $\eta_{nm}^\pm = \check{\Psi}_{j,\pm}(\zeta_{nm}^\pm)$ establishes the second assertion of the theorem. $\square$

Let us make the case $j = 0$ more explicit. Then $P_0 = 1$, $M_0 \cong \mathcal{O}(S_{qr}^2)$, and $\pi_0 \cong \tilde{\pi}_0$ is just the Heisenberg representation of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$. Accordingly, $\tilde{\pi}_0(f)\eta = f \triangleright \eta$ for $f \in \mathcal{U}_q(\mathfrak{su}_2)$, $\eta \in \mathcal{O}(S_{qr}^2)$. The 1×1 -matrices m_0 , $m_0^\dagger$ and n_0 from Subsection 6.2 have the entries $q^{-1}B$, qB^* and qA , respectively. Hence we obtain for $z, \varphi \in \mathcal{F}_b(S_{qr}^2)$

$$\tilde{\pi}_0^b(z)\varphi = z\varphi, \quad \tilde{\pi}_0^b(X)\varphi = q^{-1}\varphi B, \quad \tilde{\pi}_0^b(X^*)\varphi = q\varphi B^*, \quad \tilde{\pi}_0^b(Y)\varphi = q\varphi A. \quad (79)$$

Inserting these formulas into (76)–(78), we recover Equations (63)–(65) which we used to define a $\mathcal{U}_q(\mathfrak{su}_2)$ -action on the operator algebra $\mathcal{L}^+(\mathcal{D})$. In particular, Equations (63)–(65) and (79) give a new description of the Heisenberg representation of $\mathcal{O}(S_{qr}^2) \rtimes \mathcal{U}_q(\mathfrak{su}_2)$ on $\mathcal{O}(S_{qr}^2) \subset \mathcal{F}_b(S_{qr}^2)$ by left and right multiplications.

References

- [1] Brzeziński, T., *Quantum homogeneous spaces as quantum quotient spaces*. J. Math. Phys. **37** (1996), 2388–2399.
- [2] Brzeziński, T. and S. Majid, *Quantum geometry of algebra factorisations and coalgebra bundles*. Commun. Math. Phys. **213** (2000), 491–521.
- [3] Dijkhuizen, M. and T. K. Koornwinder, *Quantum homogeneous spaces, quantum duality and quantum 2-spheres*. Geometriae Dedicata **52** (1994), 291–315.
- [4] Fiore, G., *On the decoupling of the homogeneous and inhomogeneous parts in inhomogeneous quantum groups*. J. Phys. A **35** (2002), 657–678.
- [5] Hajac, P. M., *Bundles over quantum sphere and noncommutative index theorem*. K-Theory **21** (1996), 141–150.
- [6] Hajac, P. M. and S. Majid, *Projective module description of the q -monopole*. Commun. Math. Phys. **206** (1999), 247–264.
- [7] Klimyk, K. A. and K. Schmüdgen, *Quantum Groups and Their Representations*. Springer, Heidelberg, 1997.
- [8] Kulish, P. P. and N. Yu. Reshetikhin, *Quantum linear problem for the sine-Gordon equation and higher representations*. Zap. Nauchn. Sem. LOMI **101** (1981), 101–110.

- [9] Lunts, V. A. and A. L. Rosenberg, *Differential operators on noncommutative rings*. Sel. math. **3** (1997), 335–359.
- [10] Mimachi, K. and M. Noumi, *Quantum 2-spheres and big q -Jacobi polynomials*. Commun. Math. Phys. **128** (1990), 521–531.
- [11] Montgomery, S., *Hopf algebras and their actions on rings*. Amer. Math. Soc., Providence, R.I., 1993.
- [12] Müller, E. F. and H.-J. Schneider, *Quantum homogeneous spaces with faithfully flat module structures*. Israel J. Math. **111** (1999), 157–190.
- [13] Podleś, P., *Quantum spheres*. Lett. Math. Phys. **14** (1987), 193–202.
- [14] Schmüdgen, K., *Unbounded Operator Algebras and Representation Theory*. Birkhäuser, Basel, 1990.
- [15] Schmüdgen, K. and E. Wagner, *Hilbert space representations of cross product algebras*. J. Funct. Anal. **200** (2003), 451–493.
- [16] Schneider, H.-J., *Principal homogeneous spaces for arbitrary Hopf algebras*. Israel J. Math. **72** (1990), 167–195.
- [17] Soibelman, Ya. S. and L. L. Vaksman, *Algebra of functions on the quantum $SU(2)$* . Funct. Anal. Appl. **22** (1988), 170–181.
- [18] Sz.-Nagy, B. and C. Foias, *Analyse harmonique des operateurs de l'espace de Hilbert*. Akademiai Kiado, Budapest, 1979.
- [19] Woronowicz, S. L., *Twisted $SU(2)$ group. An example of a non-commutative differential calculus*. Publ. RIMS Kyoto Univ. **23** (1987), 117–181.